

PROPERTIES OF SATURN KILOMETRIC RADIATION MEASURED WITHIN ITS SOURCE REGION

L. Lamy^{*}, P. Schippers[†], P. Zarka^{*}, B. Cecconi^{*}, C. S. Arridge[‡],
M. K. Dougherty[§], P. Louarn[¶], N. André[¶], W. S. Kurth[†],
R. Mutel[†], D. A. Gurnett[†], and A. J. Coates[‡]

Abstract

On 17 October 2008, the Cassini spacecraft crossed the southern sources of Saturn kilometric radiation (SKR), while flying along high-latitude nightside magnetic field lines. In situ measurements allowed us to characterize for the first time the source region of an extra-terrestrial auroral radio emission. Using radio, magnetic field and particle observations, we characterize the plasma environment surrounding the SKR sources. Magnetic field lines supporting local as well as distant radio sources (detected on both extraordinary and ordinary modes) map a continuous, high-latitude and spiral-shaped unusual auroral oval observed on the dawnside, consistent with enhanced auroral activity. While investigating the Cyclotron Maser Instability (CMI) as a mechanism responsible for SKR generation, we find that observed cutoff frequencies are consistent with radio waves amplified perpendicular to the magnetic field by hot (6 to 9 keV) resonant electrons, measured locally. Finally, we confirm and quantify previous observations of the SKR elliptical polarization, found to evolve toward circular polarization with the distance to the source.

^{*} LESIA, Observatoire de Paris, UMR CNRS 8109, 92195 Meudon, France

[†] Department of Physics and Astronomy, University of Iowa, Iowa City, IA 52242, USA

[‡] Mullard Space Science Lab., University College London, London RH5 6NT, UK

[§] Blackett Lab., Imperial College, Prince Consort Road, London SW7 2BZ, UK

[¶] CESR, Université Paul Sabatier, CNRS, 31028 Toulouse 4, France

