

Literaturverzeichnis

- Adriano, D. C. 1986. Trace elements in the terrestrial environment. Springer. N.Y
- Berg, T., Steinnes, E. 1997. Recent trends in atmospheric deposition of trace elements in Norway as evident from the 1995 moss survey. *Science of the Total Environment*, 208:197-206.
- Brown, D. H. 1984. Uptake of mineral elements and their use in pollution monitoring. In Dyer, A. F., Duckett, J. G. (eds.) : The experimental biology of bryophytes. Academic Press. London. pp. 55-62.
- Burton, M.A.S. 1990. Terrestrial and aquatic bryophytes as monitors of environmental contaminants in urban and industrial areas. *Botanical Journal of the Linnean Society* 104: 267–280.
- Comeau G., Leblanc F. 1971. Influence de l'ozone et de l'anhydride sulfureux sur la regeneration des feuilles de *Funaria hygrometrica*. *Hedw. Nat. Can. (Que.)* 98: 347–358.
- Düll R. 1991. Zeigerwerte von Moosen. In: Ellenberg H. (Hrsg.): Zeigerwerte von Pflanzen in Mitteleuropa. Scripta Geobot. 18. 258pp.
- Frahm, J.-P. 1998. Moose als Bioindikatoren. Quelle u. Mayer, Wiesbaden.
- Frahm J.-P., Frey W. 1992. Moosflora. 3.Aufl. Ulmer, Stuttgart.
- Franzen I. 2001. Epiphytische Moose und Flechten als Bioindikatoren der Luftqualität am Westrand des Ruhrgebietes. *Limprichtia* 19: 1–85 + Karten.
- Gjengedal, E., Steinnes, E. 1990. Uptake of metal ions in moss from artificial precipitation. *Environmental Monitoring and Assessment* 14 : 77-87.
- Gilbert O.L. 1970. Bryophytes as indicators of air pollution in the Tyne valley. *New Phytologist* 67: 15–30.
- Gilbert O.L. 1992. Lichen reinvasion with declining air pollution. In: Bates J., Farmer A. 1992. (eds.): *Bryophytes and lichens in a changing environment*. Clarendon Press. Oxford. pp. 159–177.
- Grims F., Köckinger H. 1999. Rote Liste gefährdeter Laubmose (Musci) Österreichs. In: Niklfeld, H. (ed.): Rote Liste gefährdeter Pflanzen Österreichs. Austria Medien Service, Wien. pp. 157–171.
- Grims F., Köckinger H., Schriebl F., Suanjak M., Zechmeister H.G., Ehrendorfer F. 1999. Die Laubmose Österreichs. Catalogus Flora Austriae, II. Teil, Bryophyten (Moose), Heft 1, Musci (Laubmose). Österr. Akademie d. Wissenschaften, Wien.
- Hackl, A., Mauschitz, G. 2001. Emissionen aus Anlagen der Österreichischen Zementindustrie III. Jahresreihe 1997-1999. Bericht. Weitra / Wien
- Harmens H 2005. Heavy metals in European mosses, survey 2005/2006. Monitoring Manual. International Cooperative programme on effects of air pollution of natural vegetation and crops. ICP Vegetation Coordination Centre. Bangor.
- Herpin, U., Siewers, U., Kreimes, K., Markert, U. 2001. Biomonitoring – evaluation and assessment of heavy metal concentrations from two German moss monitoring surveys. In: Burga, C. A., Kratochwil A. (eds.) *Biomonitoring: Generals and applied aspects on regional and global scales*. Kluwer, Dordrecht, pp.73-95.
- Hill M.O., Roy D.B., Mountford J.O., Bunce R.G.H. 2000. Extending Ellenberg's indicator values to a new area: an algorithmic approach. *Journal of Applied Ecology* 37: 3–15.
- Holoubek I, Kořinek P, Šeda Z, Schneiderová E, Holoubková I, Tříška J, Cudlin P, Časlavsky J (2000): The use of mosses and pine needles to detect persistent organic pollutants at local and regional scales. *Environ Pollut* 109, 283-292
- Höfer, F., 1887. Beitrag zur Kryptogamenflora von Nieder-Österreich. *Verh. Zool.-Bot. Ges. in Österreich* 37: 379-380.

- Hohenwallner D., Zechmeister H.G., Gottfried M., Moser D., Pauli H., Reiter K., Grabherr G. 2006. Bryophytes as indicators for climate change at the alpine nival ecotone. *Journal of Applied Vegetation Science*. submitted
- Humer-Hochwimmer, K., Zechmeister, H.G. 2001. Die epiphytischen Moose im Wienerwald auf Wiener Stadtgebiet und ihre Bedeutung für die Bioindikation von Luftschadstoffen. *Limprichtia* 18: 1-99 + Anhang.
- LeBlanc F., DeSloover J., 1970. Relation between industrialization and the distribution and growth of epiphytic lichens and mosses in Montreal. *Canadian Journal of Botany* 48: 1485–1496.
- Mägdefrau, K. 1982, ‘Life-forms of bryophytes’, In: Smith, A. J. E., (ed.): ‘Bryophyte ecology’, Chapman and Hall, London. pp. 45-58.
- Manual for Integrated Monitoring. Draft 2.0. 1998. UN ECE Convention on long-range transboundary air pollution. IM Programme Centre.
- Markert, B., Breure, T., Zechmeister, H. G. 2003. Definitions, strategies and principles for bioindication / biomonitoring the environment. In: Breure, A.M., Markert, B.A., Zechmeister H.G. (eds.): Bioindicators / Biomonitoring (principles, assessment, concepts). Elsevier. Amsterdam.
- Onianwa, P.C. 2001b. Monitoring atmospheric metal pollution: a review of the use of mosses as indicators. *Environmental Monitoring and Assessment* 71: 13-50.
- Onno, M., 1941. Vegetationsreste und ursprüngliche Pflanzendecke des westlichen Wiener Stadtgebietes. *Feddes Report, Beihefte* 126: 53-127.
- Orliński R (2002): Multipoint moss passive samplers assessment of urban airborne polycyclic aromatic hydrocarbons: concentrations profile and distribution along Warsaw main streets. *Chemosphere* 48: 181-186
- Ötvös E, Kozák IO, Fekete J, Sharma VK, Tuba Z (2004): Atmospheric deposition of polycyclic aromatic hydrocarbons (PAHs) in mosses (*Hypnum cupressiforme*) in Hungary. *The Sci Total Environ* 330: 89-99
- Ramadan, Z., Song, X. H., Hopke, P. K. 2000 Identification of sources of Phoenix aerosol by positive matrix factorization. *Journal of the Air & Waste Management Association*, 50: 1308-1320
- Rao D.N. 1982. Responses of bryophytes to air pollution. In: Smith A.J.E., (ed.) *Bryophyte Ecology*. Chapman and Hall, London, pp. 445–471.
- Reiter, B., Stroh, R. 1995. Behandlung von Abfällen in der Zementindustrie. Monographien des Umweltbundesamtes 72: 1-146. Umweltbundesamt. Wien
- Saukel J., Köckinger H. 1999. Rote Liste gefährdeter Lebermoose (Hepaticae) und Hornmoose (Anthocerotae) Österreichs. In: Niklfeld, H. (ed.): Rote Liste gefährdeter Pflanzen Österreichs. Austria Medien Service, Wien, pp. 172–177.
- Schaffers A.P., Sykora K.V. 2000. Reliability of Ellenberg indicator values for moisture, nitrogen and soil reaction: a comparison with field measurements. *Journal of Vegetation Science* 11: 225–244.
- Stapper N., Franzen I., Gohrbandt S., Frahm J.P. 2000. Epiphytische Moose und Flechten kehren ins Ruhrgebiet zurück. *LÖBF-Mitteilungen* 2/2000: 12-21.
- Streit, B. 1994. Lexikon Ökotoxikologie. VCH. Weinheim
- Takeda, K., Marumoto, K., Minamikawa, T. Sakugawa, H., Fujiwara, K. 2000. Three-year determination of trace metals and the lead isotope ratio in rain and snow depositions collected in Higashi-Hiroshima, Japan. *Atmospheric Environment*, 34: 4525-4535
- Taylor, F.G, Whitterspoon, J. P. 1972. Retention of simulated fallout particles by lichens and mosses. *Health Physics* 23: 867-869.

- Umweltkontrollbericht. 2001. Umweltsituation in Österreich. 6. Bericht. UBA Wien
- Verry, E.S., Vermette, S.J. (eds.). 1992. The deposition and fate of trace metals in our environment. USDA-Forest service. MN.
- Winiwarter, W., Schneider, M. 1995. Abschätzung der Schwermetallemissionen in Österreich. Reports UBA 95-108. Umweltbundesamt. Wien
- Zechmeister, H. G., 1994. Biomonitoring der Schwermetalldepositionen mittels Moosen in Österreich. Monographien des Umweltbundesamtes Wien 42:1-168
- Zechmeister, H. G. 1995. Growth rates of five pleurocarpous moss species under various climatic conditions. *Journal of Bryology* 18: 455-468.
- Zechmeister, H. G. 1998. Annual growth of four pleurocarpous moss species and their applicability for biomonitoring heavy metals. *Environmental Monitoring and Assessment* 52: 441-451.
- Zechmeister H.G., Dirnböck, T., Hülber, K., Mirtl, M. 2006a. Assessing airborne pollution effects by bryophytes – Lessons learned by long-term integrated monitoring in Austria. *Environmental Pollution*.
- Zechmeister, H.G., Grodzinska, K., Zarek-Lukaszewska, G. 2003. Bryophytes. In: Markert, B.A, Breure, A.M., Zechmeister, H.G. (eds.): Bioindicators / Biomonitoring (principles, assessment, concepts). Elsevier. Amsterdam. pp. 329-375.
- Zechmeister, H.G., Hohenwallner, D. 2006. A comparison of biomonitoring methods for the estimation of atmospheric pollutants in an industrial town in Austria. *Environmental Monitoring and Assessment*. In press.
- Zechmeister, H.G., Hohenwallner, D., Riss, A., Hanus-Illnar, A. 2002a. Variations in heavy metal concentrations in the moss species *Abietinella abietina* (Hedw.) Fleisch. according to sampling time, within site variability and increase in biomass. *Science of the Total Environment* 301: 55-65.
- Zechmeister, H.G., Hohenwallner, D., Riss, A., Hanus-Illnar, A. 2005. Estimation of element deposition deriving from road traffic sources by mosses. *Environmental Pollution* 138: 238 - 249.
- Zechmeister, H.G., Hohenwallner, D., Hanus-Illnar, A. Roder, I., Riss, A. 2006b. Biomonitoring der Schwermetalldepositionen mittels Moosen in Österreich. Aufsammlung 2005. Report. Umweltbundesamt Wien.
- Zechmeister, H.G., Hohenwallner, D., Riss, A., Hanus-Illnar, A., Scharf., S. 2006c. Road traffic pollution measured by using mosses in a tunnel experiment in Vienna , Austria. *Environmental Science and Pollution Research*. DOI: <http://dx.doi.org/10.1065/espr2006.01.292>
- Zechmeister, H.G., Tribsch, A., Hohenwallner, D. 2002b. Die Moosflora von Linz und ihre Bedeutung für die Bioindikation. *Naturkundliches Jahrbuch der Stadt Linz* 48: 111-191.