



zu den Seiten 519–529

1: *Acanthocardia (Acanthocardia) clavata* (HILBER, 1879); Pöls bei Wildon [Holotypus zu HILBER 1879b: Taf. 6/8a+b; NHMWien 1861/I/313], x 2. – 2a+b: detto; Grund [Abb.-Orig. zu SIEBER 1956a: Taf. 2/1e + 6b; NHMWien 1963/488]. – 3a+b + 4a+b: detto; Grund [Abb.-Orig. zu SIEBER 1956a: Taf. 2/5c+d; NHMWien 1963/493], x 2. – 5: *Acanthocardia (Acanthocardia) michelottiana* (HÖRNES, 1862); Gauderndorf [Paratypus zu HÖRNES 1862; Abb.-Orig. zu SCHAFFER 1910c: Taf. 29/6 und zu STEININGER & al. 1971: Taf. 39/8; NHMWien 1860/L/426]. – 6: *Acanthocardia (Acanthocardia) mioechinata* (SCHAFFER, 1910); Gauderndorf [Syntypus zu SCHAFFER 1910c: Taf. 29/17; NHMWien 1860/L/388]. – 7a+b: *Acanthocardia (Acanthocardia) michelottiana* (HÖRNES, 1862); Gauderndorf [? Holotypus zu HÖRNES 1862: Taf. 27/4a+b; Abb.-Orig. zu SCHAFFER 1910c: Taf. 29/8; NHMWien 1860/L/426], a - x 1,5; b - x 3. – 8a+b: *Acanthocardia (Acanthocardia) moeshana* (HÖRNES, 1862); Gauderndorf [Holotypus zu HÖRNES 1862: Taf. 30/5a+b; Abb.-Orig. zu SCHAFFER 1910c: Taf. 29/10 + 11 und zu STEININGER & al. 1971: Taf. 39/4; NHMWien 1851/VI/42]. – 9: *Cardium (Bucardium) ringens danubianum* MAYER, 1866; Enzesfeld [Abb.-Orig zu HÖRNES 1862: Taf. 26/4+5; NHMWien 1853/XVI/43].




zu den Seiten 530–545

1a+b: *Acanthocardia (Acanthocardia) rugosicostata* (SCHAFFER, 1910); Gauderndorf [Holotypus zu SCHAFFER 1910c: Taf. 29/13; Abb.-Orig. zu STEININGER & al. 1971: Taf. 39/5; NHMWien 1863/XV/1259]. – 2a+b - 4: *Acanthocardia (Acanthocardia) turonica* (HÖRNES, 1862); Wien 18., Pötzelinsdorf [Syntypen zu HÖRNES 1862: Taf. 27/3a-e; NHMWien 1862/I/493]. – 5a+b + 6a+b: *Acanthocardia (Acanthocardia) vidali ritzingensis* (SIEBER, 1956); Ritzing [Holotypus zu SIEBER 1956a: Taf. 2/3a+b sowie Paratypus; NHMWien 1963/489]. – 7a+b: *Parvicardium minimum* (PHILIPPI, 1836); Vöslau [Abb.-Orig. zu SIEBER 1956a: Taf. 2/10a+b, und zu STEININGER in STEININGER, SCHULTZ, STOJASPAL & al. 1978: Taf. 15/4; NHMWien 1963/502], x 3. – 8a+b: *Parvicardium subhispidum* (HILBER, 1882); Gainfarn [Abb.-Orig. zu SIEBER 1956a: Taf. 2/8b; NHMWien 1963/503], x 4. – 9a+b: *Plagiocardium (Plagiocardium) hirsutum* (BRONN, 1831); Steinebrunn [Abb.-Orig. zu HÖRNES 1862: Taf. 26/6-9; NHMWien 1937/II/353], x 3. – 10a+b: *Plagiocardium (Papillocardium) papillosum* (POLI, 1791); Wien 16., Ottakring [Abb.-Orig. zu SIEBER 1956a: Taf. 1/16, und zu STEININGER in STEININGER, SCHULTZ, STOJASPAL & al. 1978: Taf. 15/3; NHMWien 1963/504], x 3. – 11a+b: *Nemocardium (Discors) aquitanicum laevinflatum* (SACCO, 1899); Forchtenau [Abb.-Orig. zu HÖRNES 1862: Taf. 24/6+7, und zu SIEBER 1956a: Taf. 3/8; NHMWien 1853/III/182].




zu den Seiten 534–538, 565

1a+b + 2a+b: *Acanthocardia (Europicardium) multicostata* (BROCCHI, 1814); Grund [Abb.-Orig. zu HÖRNES 1862: Taf. 30/7a-c, zu SIEBER 1956a: Taf. 3/10a+b; NHMWien 1851/XXVI/58]. – 3: *Acanthocardia (Rudicardium) grandis tereticostalis* (STEININGER, 1963); Fels am Wagram [Holotypus zu STEININGER 1963b: Taf. 9/1a+b; Abb.-Orig. zu STEININGER & al. 1971: Taf. 41/1+4; PIWien]. – 4a+b: *Acanthocardia (Rudicardium) grandis* (HÖLZL, 1958); Fels am Wagram [Abb.-Orig. zu STEININGER 1963b: Taf. 8/1a+b; und zu STEININGER & al. 1971: Taf. 39/7, Taf. 40/1; PIWien]. – 5a+b: *Limnopagetia moravica moravica* (RZEHAK, 1883); Oslavany, Tschechische Republik [nicht abgebildeter Paratypus zu *brusinai* RZEHAK 1893 und Abb.-Orig. zu CTYROKY 1972: Taf. 9/7+8; NHM Wien 1966/659/B], x 2. – 6a+b: detto [nicht abgebildeter Paratypus zu RZEHAK 1893 (vgl. CTYROKY 1972: 96); NHM Wien 1966/656/C], x 2. – 7a+b: detto [Lectotypus zu CTYROKY 1972: Taf. 8/2-5; Abb.-Orig. zu CTYROKY in STEININGER & al. 1973: Taf. 24/1a+b; NHM Wien 1966/656/A], x 2.



zu den Seiten 545–550

1a+b: *Laevicardium (Laevicardium) sandbergeri* (GÜMBEL, 1861); Fels am Wagram [Abb.-Orig. zu STEININGER 1963b: Taf. 9/3; und zu STEININGER & al. 1971: Taf. 37/2; PIWien]. – 2 + 3: *Nemocardium (Discors) spondyloides* (HAUER, 1847); Steinebrunn [Abb.-Orig. zu HÖRNES 1862: Taf. 24/1-3; GBAWien. – Innenansichten zu Taf. 80/2+3].



zu den Seiten 545–547

1a+b: *Nemocardium (Discors) spondyloides* (HAUER, 1847); Enzesfeld [Abb.-Orig. zu HÖRNES 1862: Taf. 24/4+5; NHMWien 1855/II/64]. – 2 + 3: *Nemocardium (Discors) spondyloides* (HAUER, 1847); Steinebrunn [Abb.-Orig. zu HÖRNES 1862: Taf. 24/1-3; GBAWien. – Außenansichten zu Taf. 79/2+3].




Tafel 81 + Tafel 82
zu den Seiten 548–550

1a+b: *Laevicardium (Laevicardium) kuebecki* (HAUER, 1847); Loibersdorf [Abb.-Orig. zu HÖRNES 1862: Taf. 21/2; zu SCHAFFER 1910c: Taf. 31/2; NHMWien 1848/III/75]. – 2a+b: *Laevicardium (Laevicardium) kuebecki* (HAUER, 1847); Loibersdorf [Abb.-Orig. zu SCHULTZ 1998: 100, Taf. 44/1a+b; NHMWien 1866/XL/16].



Tafel 83




zu den Seiten 548–553, 562–563

1a+b: *Laevicardium (Habecardium) subturgidum* (d'ORBIGNY, 1852); Steinebrunn [NHMWien 1857/XIV/38]. – 2a+b: detto; Gainfarn [NHMWien 1863/XV/1285]. – 3a+b: *Laevicardium (Laevicardium) kuebecki* (HAUER, 1847); Loibersdorf [NHMWien 1866/XL/18], a - x 1; b, Detailansicht x 1,5. – 4: ? *Korobkoviella lipoldi* (ROLLE, 1858); Motschnik-Mühle im Liffaygraben, bei Prasberg [Slowenien; Syntypus zu ROLLE 1859: Taf. 2/10a; NHMWien 1858/XIV/42], x 2. – 5: detto [Syntypus zu ROLLE 1859; NHMWien 1858/XIV/41], x 3. – 6a+b: *Korobkoviella cf. kiktentoi* (MERKLIN, 1974); Ottenthal, Untere Leithen [Abb.-Orig. zu BALDI in SEIFERT, BRAUNSTEIN & BALDI: Taf. 4/1, rev. POPOV, 29.IV.1999; PIWien], x 4.

**zu den Seiten 554–555**

1a+b + 3: *Laevicardium (Habecardium) tenuisulcatum* (Nystr., 1836); Loibersdorf [Abb.-Orig. zu HÖRNES 1862: Taf. 25/1a-d; zu SCHAFFER 1910c: Taf. 29/14, Taf. 30/1; und zu STEININGER & al. 1971: Taf. 21/3; NHMWien 1852/XII/6]. – 2: detto; Loibersdorf [Abb.-Orig. zu SCHAFFER 1910c: Taf. 29/15; NHMWien 1852/XII/5].

Tafel 85



zu den Seiten 557–562, 582–583, 599–600

1a+b: *Cerastoderma arcellum* (DUJARDIN, 1837); ? Niederkreuzstetten [? Abb.-Orig. zu HÖRNES 1862: Taf. 25/3; NHMWien 2003z0016/0001], a - 1:1, b - x 1.5. – 2a+b: detto; Niederkreuzstetten [NHMWien 1859/XLV/89], x 2. – 3a-c: detto; Laa/Thaya [NHMWien 1860/L/139], a - 1:1, b+c - x 2. – 4+5: *Cerastoderma edule commune* (MAYER, 1866); Gauderndorf [Abb.-Orig. zu STEININGER & al. 1971: Taf. 37/7; GBAWien]. – 6: detto; Gauderndorf [Belegmaterial zu HÖRNES 1862; NHMWien 1851/VI/43]. – 7a+b + 8a+b: *Cerastoderma edule felsense* (STEININGER, 1963); Fels am Wagram [Holo- und Paratypus zu STEININGER 1963b: Taf. 8/2-5; Abb.-Orig. zu STEININGER & al. 1971: Taf. 39/2+3; PIWien 1660], x 1.5. – 9a+b: *Cerastoderma edule greseri* (WOLFF, 1897); Fels am Wagram [Abb.-Orig. zu STEININGER 1963b: Taf. 8/6; und zu STEININGER & al. 1971: Taf. 39/1; PIWien]. – 10a+b: *Cerastoderma edule* indet. subspec. 1; Grund [Abb.-Orig. zu SIEBER 1956a: Taf. 2/7; NHMWien 1963/498]. – 11 + 12: *Cardium carasi* (JEKELIUS, 1944); Bad Gleichenberg [Holotypus zu *Cardium gleichenbergense* PAPP, 1954: Taf. 12/12a+b; Abb.-Orig. zu PAPP 1974c: Taf. 13/13+14; NHMWien 1329/1970, Koll. PAPP]. – 13a+b: *Planocardium doengingki praeforme* (PAPP, 1954); Wiesen B [Holotypus zu PAPP 1954: Taf. 13/9; Abb.-Orig. zu PAPP 1974c: Taf. 14/14; NHMWien 1329/1970, Koll. PAPP], x 2. – 14a+b + 15a+b: *Obsoletiforma (Obsoletiforma) ghergutai* (JEKELIUS, 1944); Naxing [Abb.-Orig. zu PAPP 1954: Taf. 13/5+6; und zu PAPP 1974c: Taf. 14/5+6; NHMWien 1329/1970, Koll. PAPP], x 1.5. – 16a+b + 17a+b: *Inaequicostata ? janoschekii* (PAPP, 1954); Siebenhirten, N Mistelbach [Holo- und Paratypus zu PAPP 1954: Taf. 15/1+2; Abb.-Orig. zu PAPP 1974c: Taf. 14/10+11; NHMWien 1329/1970, Koll. PAPP], x 4. – 18a+b: detto; Waldhof bei Graz [det. PAPP; NHMWien grün A2332], x 5.