**universität
wien**

University of Vienna

Faculty of Life Science

Department of Botany and Biodiversity Research

Division of Conservation Biology,

Vegetation Ecology and Landscape Ecology

in cooperation with

**University of Natural Resources
and Life Sciences, Vienna**

Department of Spatial-, Landscape-
and Infrastructure- Sciences

Institute of Landscape Development,
Recreation and Conservation Planning

Funded by

Earth System Science

Man and
the Biosphere
Programme

ÖAW

AUSTRIAN
ACADEMY OF
SCIENCES

Supported by
Biosphärenpark

Wienerwald

Management GmbH

Lebensregion
Biosphärenpark
Wienerwald

Biosphere Reserve Integrated Monitoring based on Social and Environmental Ecosystem Services – BRIMSEN

FINAL REPORT

ISBN 978-3-7001-8617-5
DOI 10.1553/MAB-BRIMSEN

Biosphere Reserve Integrated Monitoring based on Social and Environmental Ecosystem Services - BRIMSEN

UNIVERSITY OF VIENNA

Faculty of Life Science

Department of Botany and Biodiversity Research

Division of Conservation Biology, Vegetation Ecology and Landscape Ecology (CVL)

1030 Vienna, Rennweg 14

T +43 1 4277-54382

Ass. -Prof. Dr. Thomas Wrбка (thomas.wrbka@univie.ac.at)

Thomas Wrбка is partner and co-ordinator of numerous research projects in the field of landscape ecology, vegetation and restoration ecology, and conservation biology. Special research topics comprise ecosystem services and green infrastructure, management of protected areas, rewilding, and biodiversity conservation in agricultural landscapes.

Mita Drius, PhD (mita.drius@univie.ac.at)

Mita Drius is an ecologist with focus on the assessment of ecosystem services. She works as post-doctoral project manager for the Department of Botany and Biodiversity Research, Division of Conservation Biology, Vegetation Ecology and Landscape Ecology at the University of Vienna.

UNIVERSITY OF NATURAL RESOURCES AND LIFE SCIENCES

Department of Landscape, Spatial and Infrastructure Sciences

Institute of Landscape Development, Recreation and Conservation Planning (ILEN)

1190 Vienna, Peter Jordan-Straße 82

T +43 1 47654-7204

Ao. Univ.Prof. DI Dr. Christiane Brandenburg (christiane.brandenburg@boku.ac.at)

Christiane Brandenburg is professor at the Institute of Landscape Development, Recreation and Conservation Planning, University of Natural Resources and Life Sciences, Vienna. She was project partner in several MAB projects. Her expertise includes participation and transdisciplinary work, while her focus in scientific research is on landscape-related recreation and cultural landscape development.

DI Katharina Sams (katharina.sams@boku.ac.at)

Katharina Sams is a landscape planner with research focus on nature-related recreation in an urban context as well as the assessment of socio-cultural ecosystem services in cultural landscapes. She works at the Institute of Landscape Development, Recreation and Conservation Planning.

Friedrich Knopper, BSc (friedrich.knopper@students.boku.ac.at)

Friedrich Knopper is a student of landscape planning. He works as assistant for GIS at the Institute of Landscape Development, Recreation and Conservation Planning and at the Institute of Hydrobiology and Aquatic Ecosystem Management.

Project period: March 2018 – May 2019

Suggested citation: Mita Drius, Katharina Theresa Sams, Friedrich Knopper, Christiane Brandenburg, and Thomas Wrбка, 2019. Biosphere Reserve Integrated Monitoring based on Social and Environmental Ecosystem Services – BRIMSEN. Final Report.

Keywords: Wienerwald Biosphere Reserve, Landscape Services, Ecosystem Services, Biosphere Reserve Integrated Monitoring

Acknowledgements

This project was funded by the Austrian Academy of Sciences, under the Earth System Science - Man and the Biosphere Programme. The authors would like to thank Mag. Dr. Günter Köck for the support over the project duration.

Our special thanks go to the staff of the Biosphärenpark Wienerwald Management GmbH, especially to the Director Dr. Herbert Greisberger and to DI Harald Brenner for the productive collaboration throughout the project. In the project, the staff of the Biosphärenpark Wienerwald Management GmbH had two roles: they acted both as facilitators by providing data and hosting meetings, and as first receivers of the project results.

The authors would like to thank the following data providers:

- Wolfgang Schranz of the Biosphärenpark Wienerwald Management GmbH, for providing the geodata of the “Biotoptypenkartierung Offenland” and of the borders of the Wienerwald Biosphere Reserve, and also for giving advice and background information on the geodata.
- The Austrian Federal Forests (ÖBf) for providing forest geodata.
- The Institute of Hydrobiology and Aquatic Ecosystem Management (IHG) of the University of Natural Resources and Life Sciences for providing geodata on the hydropower plants.
- The Wienerwald Tourismus GmbH for providing geodata on the recreational trails.

The authors are very grateful to Mag. Dr. Christa Hainz-Renetzeder, supervisor of the lecture “GIS-gestütztes Projekt” in the fall semester 2018/2019 at the University of Natural Resources and Life Sciences, and to her students Lena Graf, Daniel Hartl, Eric Kraus, Michele Misztowt, Tobias Riffel and Theresa Rockenschaub for providing three case studies located in the Wienerwald Biosphere Reserve and focussed on the spatial analysis of landscape services.

Preface

BRIMSEN take roots on an extensive experience gained over years on the assessment of ecosystem and landscape services, such as in the project TransEcoNet “Transnational Ecological Networks” (Central Europe programme), where the potential and the current status of ecosystem services were identified for the region of Neusiedler See, and in the MAB projects BIOSERV “Biodiversity and Ecosystem Service as scientific foundation for the sustainable implementation of the Redesigned Biosphere Reserve Neusiedler See” and CCR “Climate Change Response of Ecosystem Services in the sensitive area of Neusiedler See-Seewinkel” where the ecosystem and landscape services approach was employed to boost the spatial planning of the region of Neusiedler See. Moreover, our team has been working in the Wienerwald Biosphere Reserve since long time, as proven by the MAB projects ISWI MAB “Integrated Sustainable Wildlife Management in the Wienerwald Biosphere Reserve”, and IESP “Towards Integrated Ecological Spatial Planning for the Wienerwald Biosphere Reserve”.

Based on the integrative assessment framework for ecosystem and landscape services resulting from our previous studies and by including new approaches, we are now able on the one hand to propose a set of indicators for a Biosphere Reserve Integrated Monitoring, and on the other hand to address scientific challenges. Our project team has in-depth expertise and access to existing data sets from the research projects „Detailplanung zum Biosphärenpark Wienerwald - Bereich Offenland (2004 - 2006)“, „Offenlanderhebung des Biosphärenparks Wienerwald (2011 - 2012)“, „Machbarkeitsstudie Wienerwald - Eignung des Wienerwaldes für einen Nationalpark oder Biosphärenpark“ and from the related publications (Reimoser et al. 2008; Reimoser et al. 2012; Grabherr et al. 1998; Arnberger et al. 2002; Bianchin et al. 2012; Hermann et al. 2014; Hainz-Renetzeder et al. 2014; Wrbka et al. 2012; Kuttner et al. 2013).

Abstract

It is nowadays globally recognized that biodiversity, i.e. the diversity of ecosystems and species, sustains human life by means of the so-called ecosystem services. Landscapes, being the product of human interaction with the environment, support well-being not only via benefits from natural and semi-natural ecosystems, but also through human-modified landscape elements. Landscape services are therefore those benefits supplied by both natural and human-modified landscape elements, and they include not only potentials, materials and processes of nature but also services of cultural elements and constructions that come into being through human creation.

Biosphere Reserves are considered as means for the people who live and work within them to attain a balanced relationship with the natural and semi-natural environment. Moreover, they contribute to the needs of society by showing a way to a more sustainable future. Being devoted to nature conservation coupled with sustainable development, they rely on integrated monitoring and sustainability assessments. Landscape services lend themselves to this purpose in an optimal way as they not only can measure ecological and socio-cultural sustainability but are also a good communication tool with stakeholders and the local population.

The Wienerwald Biosphere Reserve partly surrounds the city of Vienna and other minor settlements, and it offers a heterogeneous landscape capable of delivering a variety of landscape services. The analysis of the services capacity offered by the open land elements of Wienerwald may represent a well-founded basis for the sustainable planning of the Reserve, and it can concretely contribute to a Biosphere Reserve Integrated Monitoring (BRIM).

In this work we quantify and map the capacity of landscape services offered by the open land of Wienerwald, and we offer a set of indicators for a BRIM. Starting from a high-resolution dataset, we aggregate the open land landscape elements into open land classes, and we select additional spatial indicators able to represent all landscape services, which cover social, cultural, ecological and productive spheres. Subsequently, we build on expert-based assessments to score each service through a capacity matrix. The landscape services scores are then spatialized, through an innovative analysis focused on the INTensity and DEnsity landscape service Capacity (INDEC). First, each landscape element is converted into a point, whose landscape services capacity is represented by an outer buffer linked to the scores from the capacity matrix and the size of the element. Then, touching or overlapping buffers are merged, obtaining dissolved clusters representing the cumulative intensity of landscape services capacity. In this way, we obtain 22 maps of landscape service capacity, which can be fruitfully adopted by spatial planners and decision makers - Biosphere Reserve Management *in primis*.

Subsequently, we select a set of suitable landscape services to act as indicators for a BRIM. The criteria used to accomplish the selection are representativeness, communication and relevance for the Biosphere Reserve Management. The representativeness is provided by landscape services addressed by a high amount of landscape service classes and therefore high capacity within the Wienerwald Biosphere Reserve. Landscape services that convey clarity and are easily understandable for the Wienerwald Biosphere Reserve Management or potentially involved stakeholders produce a set of indicators useful also for communication purposes. Additionally, we emphasise landscape services with an overarching character, but

we also consider a balance between ecological and socio-cultural aspects, as both are very relevant for the Wienerwald Biosphere Reserve Management. The eight landscape services are Local climate and air regulation, Refugium, Soil retention, Water supply, Recreation, Aesthetic information, Science and Education, and Food.

This study provides an innovative method for mapping landscape service capacity: unlike other studies, all landscape elements, from meadows to field crosses to roads, are treated at the same level, since the INDEC is substantially based on their location and proximity, and on the capacity matrix scores. The method allowed an appropriate balance among isolated and large landscape elements (e.g. cultivated field, meadow) and small-scale but numerous landscape elements, such as tree rows and benches. The study also highlights advantages and disadvantages of the expert-based assessment for scoring the landscape service capacity.

The capacity maps showed a great variability in the provision of service capacity. Some services were highly represented in the Biosphere Reserve (e.g. Soil formation and Nutrient Regulation), while others were underrepresented (e.g. Raw Materials). The variability evidenced through the maps is discussed in light both of the approach followed and of the data accuracy.

From a management perspective, the study offers valuable insights for an integrated monitoring of the sustainability performance of the Reserve through a thoughtful set of landscape services. Moreover, by focussing on the open land areas of the Biosphere Reserve and by excluding the forest, high value to the multifunctionality and connectivity of small-scale elements is provided. The landscape service maps can be fruitfully adopted by the Reserve Management to identify hot and cold spots for certain service, and to orient their measures accordingly.

Table of Contents

1	Introduction and aims	12
1.1	Central focus.....	12
1.2	Project aims	13
1.3	Methodological approach	13
1.4	Target groups of the results	13
2	Theoretical background	15
2.1	The concept of sustainable landscape development and the role of biosphere reserves	15
2.2	The Biosphere Reserve Integrated Monitoring	16
2.3	The concept of ecosystem services and of landscape services	17
2.4	Service capacity and capacity matrix.....	19
3	Research questions	21
4	Material and Methods	22
4.1	Project area description	22
4.2	Data sources	23
4.3	Definition of landscape services addressed in the project	24
4.4	Landscape elements classification.....	25
4.4.1	Open land habitats	26
4.4.2	Additional spatial indicators	27
4.5	Capacity matrix: assessing service capacities	27
4.6	Spatial data preparation.....	29
4.7	Transformation of spatial data	29
4.8	INDEC application.....	33
4.9	Refinements to the methodology	35
4.9.1	Exclusion of landscape services	35
4.9.2	Validation of plausibility of the landscape services capacity maps	35
4.9.3	Weighing factor adjustment	36
4.9.4	Treatment of polylines.....	36
4.9.5	Validation of the capacity matrix.....	37
4.10	Selection process of landscape services for BRIM.....	37
5	Results	39
5.1	Capacity matrix.....	39
5.2	Landscape service capacity maps	52
5.2.1	Local climate and air regulation.....	53
5.2.2	Disturbance prevention	55
5.2.3	Water regulation	57
5.2.4	Water supply.....	59
5.2.5	Soil retention	61
5.2.6	Soil formation	63
5.2.7	Nutrient regulation	65
5.2.8	Pollination.....	67
5.2.9	Refugium.....	69
5.2.10	Nursery	71
5.2.11	Food	73
5.2.12	Raw materials	75
5.2.13	Genetic resources	77
5.2.14	Medicinal resources.....	79
5.2.15	Cultivation.....	81
5.2.16	Transportation	83
5.2.17	Tourism facilities.....	85
5.2.18	Aesthetic information	87
5.2.19	Recreation.....	89
5.2.20	Cultural and artistic information	91
5.2.21	Spiritual and historic information	93
5.2.22	Science and education	95

5.3	Selected landscape services for a BRIM	97
6	Discussion.....	99
6.1	Integrated discussion on the landscape services capacity in the Biosphere Reserve	99
6.2	Considerations on data availability and data treatment	101
6.2.1	Data sources	101
6.2.2	Data classification	102
6.3	Reflections on the methodology	103
6.3.1	The capacity matrix: advantages and disadvantages	103
6.3.2	INtensity and Density of service Capacity: advantages and disadvantages	104
6.4	Management implication	105
7	Conclusions	106
8	References.....	108
9	Attachments.....	115
9.1	Definition of landscape services	115
9.2	Additional Spatial Indicators sources	118
9.3	Reclassification of open land elements	119
9.4	Classification of Additional Spatial Indicators	123
10	Teaching Landscape Services in GIS - three case studies	124

List of figures

Figure 1: From observation to monitoring	17
Figure 2: Conceptual model showing relations of ecosystem functions, services and benefits.....	19
Figure 3: Study area: the open land of the Wienerwald Biosphere Reserve.	23
Figure 4: Scheme of the approach for comparing the CICES and BRIMSEN classifications.	25
Figure 5: Example of open land classification	26
Figure 6: Workflow of the INDEC calculation, with data preparation and INDEC application.....	30
Figure 7: Scheme for point distance calculation	31
Figure 8: Graph exemplifying the theoretical assumption for defining the maximum buffer size.....	32
Figure 9: Calculation of weighting factor based on relative part.....	34
Figure 10: Process of landscape services selection for a BRIM.....	38
Figure 11: Local climate and air regulation capacity in the Wienerwald Biosphere Reserve.	53
Figure 12: Disturbance prevention capacity in the Wienerwald Biosphere Reserve.....	55
Figure 13: Water regulation capacity in the Wienerwald Biosphere Reserve.	57
Figure 14: Water supply capacity in the Wienerwald Biosphere Reserve.	59
Figure 15: Soil retention capacity in the Wienerwald Biosphere Reserve.....	61
Figure 16: Soil formation capacity in the Wienerwald Biosphere Reserve.....	63
Figure 17: Nutrient regulation capacity in the Wienerwald Biosphere Reserve.....	65
Figure 18: Pollination capacity in the Wienerwald Biosphere Reserve.	67
Figure 19: Refugium capacity in the Wienerwald Biosphere Reserve.	69
Figure 20: Nursery capacity in the Wienerwald Biosphere Reserve.....	71
Figure 21: Food capacity in the Wienerwald Biosphere Reserve.....	73
Figure 22: Raw materials capacity in the Wienerwald Biosphere Reserve.....	75
Figure 23: Genetic resources capacity in the Wienerwald Biosphere Reserve.....	77
Figure 24: Medicinal resources capacity in the Wienerwald Biosphere Reserve.	79
Figure 25: Cultivation capacity in the Wienerwald Biosphere Reserve.	81
Figure 26: Transportation capacity in the Wienerwald Biosphere Reserve.....	83
Figure 27: Tourism facilities capacity in the Wienerwald Biosphere Reserve.	85
Figure 28: Aesthetic information capacity in the Wienerwald Biosphere Reserve.	87
Figure 29: Recreation capacity in the Wienerwald Biosphere Reserve.	89
Figure 30: Cultural and artistic information capacity in the Wienerwald Biosphere Reserve.....	91
Figure 31: Spiritual and historic information capacity in the Wienerwald Biosphere Reserve.	93
Figure 32: Science and education capacity in the Wienerwald Biosphere Reserve.....	95
Figure 33: Scheme showing the spheres of the BR environment.	97
Figure 34: Landscape services for a BRIM for the Wienerwald Biosphere Reserve.....	98
Figure 35: Comparison between the inclusion and exclusion of the forest.....	102

List of tables

Table 1: Extract of BRIMSEN capacity matrix.	28
Table 2: Width distribution factor applied to polyline landscape elements (LE).....	36
Table 3: Capacity matrix of the open land (OL) classes 1-11	40
Table 4: Capacity matrix of the open land (OL) classes 12-24	42
Table 5: Capacity matrix of the open land (OL) classes 25-36	44
Table 6: Capacity matrix of open land (OL) classes 37-50.....	46
Table 7: Capacity matrix of the open land (OL) classes 51-62	48
Table 8: Capacity matrix of the Additional Spatial Indicators classes 1-18.....	50
Table 9: Three clusters with highest INDEC for Local climate and air regulation.	54
Table 10: Three clusters with highest INDEC for Disturbance prevention.....	56
Table 11: Three clusters with highest INDEC for Water regulation.	58
Table 12: Three clusters with highest INDEC for Water supply.	60
Table 13: Three clusters with highest INDEC for Soil retention.	62
Table 14: Three clusters with highest INDEC for Soil formation.....	64
Table 15: Three clusters with highest INDEC for Nutrient regulation.....	66
Table 16: Three clusters with highest INDEC for Pollination.....	68
Table 17: Three clusters with highest INDEC for Refugium.	70
Table 18: Three clusters with highest INDEC for Nursery.	72
Table 19: Three clusters with highest INDEC for Food.....	74
Table 20: Three clusters with highest INDEC for Raw materials.	76
Table 21: Three clusters with highest INDEC for Genetic resources.....	78
Table 22: Three clusters with highest INDEC for Medicinal resources.	80
Table 23: Three clusters with highest INDEC for Cultivation.	82
Table 24: Three clusters with highest INDEC for Transportation.....	84
Table 25: Three clusters with highest INDEC for Tourism facilities.	86
Table 26: Three clusters with highest INDEC for Aesthetic information.....	88
Table 27: Three clusters with highest INDEC for Recreation.	90
Table 28: Three clusters with highest INDEC for Cultural and artistic information.	92
Table 29: Three clusters with highest INDEC for Spiritual and historic information.	94
Table 30: Three clusters with highest INDEC for Science and education.....	96
 Table A 1: BRIMSEN's definitions of the landscape services.....	 115
Table A 2: Data sources of the Additional Spatial Indicators (ASI) classes,.....	118
Table A 3: Reclassification of the open land types.....	119
Table A 4: Classification of Additional Spatial Indicators	123

Abbreviations

ASI	Additional Spatial Indicators
BR	Biosphere Reserve
BRIM	Biosphere Reserve Integrated Monitoring
INDEC	Analysis based on INtensity and DEnsity of landscape service Capacity
LE	Landscape Element
LS	Landscape Service
OL	Open Land

1 Introduction and aims

The project BRIMSEN “Biosphere Reserve Integrated Monitoring based on Social and Environmental Ecosystem Services” aimed at analysing the ecosystem services occurring in the Wienerwald Biosphere Reserve, so to produce regional maps of service capacity distribution, and to contribute a selection of indicators for a Biosphere Reserve Integrated Monitoring (BRIM).

1.1 Central focus

The central theme of the project is the analysis of the link between natural and semi-natural ecosystems and human well-being, grouped under the concept of ecosystem services.

The publication of the Millennium Ecosystem Assessment (MEA 2005) has stimulated widespread, international debate on the importance of the links between ecosystems and human well-being. However, benefits from ecosystems and from landscapes are often not fully taken into account in environmental planning and decision-making. Based on incomplete information, multi-functional, productive landscapes are used in an unsustainable way, fact that leads to the degradation of our “natural capital” at the expense of our own welfare and that of future generations (TEEB 2010). Therefore, as it was postulated in Nagoya 2010 as well as in the European Biodiversity Strategy to 2020, our main future challenges are to stop the loss of biodiversity and to reduce the global impact on our natural resources.

The conceptual framework adopted in the study is a conceptual variation of ecosystem services, i.e. landscape services. Landscapes are spatially heterogeneous geographic areas characterized by diverse interacting patches or ecosystems, ranging from relatively natural terrestrial and aquatic systems such as forests, grasslands, and lakes to human-dominated environments including agricultural and urban settings (Forman 1995). Since a landscape can provide services to the society through its multi-functionality and the processes that emerge from a set of ecosystems, the so-called landscape services are those benefits delivered by both natural ecosystems and man-made landscape elements (de Groot 2006; Konkoly-Gyuró 2011). Landscape services can provide insights on the current state of land management, providing a way to monitor its sustainability over time, when adopted as indicators (Termorshuizen et al. 2009).

The study is set in the Wienerwald Biosphere Reserve. Biosphere reserves are seen as model regions for nature conservation practices reconciled with sustainable socio-economic development, and as such they are well suited for “living laboratories” for studying, testing out and demonstrating integrated management of land, water and biodiversity in a sound balance with human uses. In BRIMSEN, we focused on two of the three pillars of sustainability, due to the project team’s expertise: they are the environmental and socio-cultural perspectives. According to the UNESCO (2005), synergies between cultural diversity and biological diversity are regarded as crucial factors for the preservation and balance of social and ecological systems.

1.2 Project aims

The main objective of the project was the development and provision of technical and methodological framework for the regional assessment and mapping of landscape services. We chose Wienerwald Biosphere Reserve as case study area.

The main objective can be divided into the following specific sub-goals:

- Define suitable landscape elements to represent landscape services in light of spatial data availability
- Simultaneously assess and map ecological and socio-cultural landscape services
- Provide a rapid and replicable toolset for landscape services assessment, that would contribute to a replicable monitoring of the Biosphere Reserve's sustainability performance
- Select landscape services as indicators for a BRIM

1.3 Methodological approach

The project's methodological approach has the following basic conceptual considerations:

- The basic spatial entity of this study is the "landscape element", in the sense of (Forman et al. 1986) and Forman (1995). Accordingly, in our study a landscape element is any spatial element in the "open" agrarian part of the Biosphere Reserve.
- All landscape elements, ranging from field crosses to meadows, roads and forest edges, are spatially treated at the same level, this way guaranteeing that all landscape services receive the same consideration.
- The ranking of each landscape service capacity is performed through a capacity matrix and built on an expert-based judgement, tuned for the study area. Each landscape element was assigned a score based on its unconnected capacity to potentially deliver a service.
- The method to spatialize the scores from the capacity matrix follows the concept of spatial connectivity. Ecological spatial connectivity refers to processes by which genes, organisms, populations, species, nutrients, and/or energy move among spatially distinct habitats, populations, communities, or ecosystems (Carr et al. 2017). In this project, the spatial connectivity of the landscape elements is based on their neighbourhood or proximity. Consequently, the spatial connectivity of each service is related both to the score assigned through the capacity matrix (intensity of the service capacity) and to the proximity among landscape elements (density of the service capacity).

1.4 Target groups of the results

The results of the project are aimed at all actors in the Biosphere Reserve who are involved in the formation in conflict potentials for different land use management options, or/and are affected by the consequences of such conflicts. Target groups therefore include for example, the Biosphärenpark Wienerwald Management GmbH, responsible decision makers in provincial and district administrations, local councils, representatives of different recreational user groups, landowners and managers, associations and advocacy groups. With regard to the sectoral or technical responsibilities, in particular the areas of leisure industry, nature and species protection, spatial planning, forestry, agriculture and wildlife

management can be addressed. Outside the project area's stakeholders, other target groups of this study are other National Parks and biosphere reserves *in primis*, located inside and outside Austria, which might replicate the methodology, tuning it to the characteristics of their region, but also governmental institutions, agencies and research institutes.

2 Theoretical background

2.1 The concept of sustainable landscape development and the role of biosphere reserves

The Man and Biosphere Programme can be considered as a UNESCO flagship programme linking nature and landscape to sustainable development. The International Coordinating Council of the UNESCO “Man and the Biosphere” (MAB) Programme convened for the first time in 1971 and laid the foundations for a new type of conservation area – Biosphere Reserves (Ruoss 2013). Today, the world network of 610 Biosphere Reserves can praise experience gathered over 40 years as model regions for sustainable development to climate change mitigation and adaptation and aims at making a substantial contribution to these processes as well as to the conservation of biological diversity (Ruoss 2013).

The principles of sustainable landscape development imply that environmental, socio-cultural and economic functions should be balanced in space and time to maintain their potential to provide goods and services to future generations (WCED 1987; Linehan et al. 1998; UNESCO 2005). However, short-term goals mainly related to economic values tend to dominate over public landscape values. To meet this challenge, environmental, socio-cultural and economic value domains have to be integrated in landscape planning processes by taking into account that each of them requires a different way of looking at the functions and pattern of landscapes at different scales (Opdam et al. 2006). Understanding these three value domains and their interrelations, adverse effects as well as synergies and trade-offs, is seen as the fundamental basis for sustainable landscape development (Pope et al. 2004).

Biosphere reserves provide an example of an integrated sustainability framework, which explicitly acknowledges that complex socio-economic and ecological systems are inextricably linked (Levrel et al. 2008). Therefore, they have a primary role in coupling nature conservation practices with sustainable socio-cultural development (Lotze-Campen et al. 2008). The so-called “Seville Strategy” was elaborated during an international conference in Seville, Spain, in 1995 and re-confirmed during the Seville +5 conference in Pamplona in 2000 (UNESCO 2002). Based on the “Seville Strategy”, ecological and socio-economic assessments are a prerequisite for spatial zoning of the biosphere reserves, reinforcing scientific research and initiating development tasks (UNESCO 1996; UNESCO 2005). Indeed, each biosphere reserve is intended to fulfil three basic complementary and mutually reinforcing functions:

- a conservation function - to contribute to the conservation of landscapes, ecosystems, species, and genetic variation;
- a development function - to foster economic and human development that is socio-culturally and ecologically sustainable;
- a logistic function - support for demonstration projects, environmental education and training, research, and monitoring related to local, national, and global issues of conservation and sustainable development. (UNESCO 2002)

Biosphere reserves are seen as models for sustainable development according to the Seville Strategy. A model in this context can be defined as “best practice” regarding integrated approach for sustainable development, following the principle of a balanced growth of the three pillars of sustainability: protecting the natural and cultural heritage and creating added

values for people. It has to aim at facilitating the reproduction of the achievements in other areas as well (Ruoss 2013).

Although the role of Biosphere Reserves as models for sustainable development was promoted by UNESCO as a key contribution to the post-Rio+20 process, some key questions which would strengthen and enhance the significance of the Biosphere Reserve as a model for Sustainable Development have not yet been fully considered in the new generation of Biosphere Reserves (Ruoss 2013). Some of these critical issues are:

- dealing with natural capital, the core idea of the green economy concept;
- enhancing awareness of values related to natural resources: biodiversity, landscape;
- increasing added value for people through sustainable growth;
- relating cultural diversity and heritage to economy and environment;
- dedicating and linking innovation and research to development;
- transferring knowledge and know-how to the general public;
- involving stakeholders and public in decision processes;
- assessing heritage protection and development in territorial processes (e.g. footprint).

These complex and multidisciplinary aspects can be simultaneously addressed through a cohesive monitoring strategy, which couples the socio-cultural with ecological spheres into an integrated monitoring. A useful tool in this sense is provided by the BRIM.

2.2 The Biosphere Reserve Integrated Monitoring

In 2003, the Austrian MAB committee has decided to devote at least part of its research funds to the development of Biosphere Reserve Integrated Monitoring (BRIM), in support of the international MAB agenda in general and Austrian biosphere reserves in particular (Fischer-Kowalski et al. 2004).

The monitoring platform BRIM operates within the World Network of Biosphere Reserves (WNBR), providing abiotic, biological, and socio-economic information. In this context, within the UNESCO MAB Biosphere Reserves Directory, one can perform queries by research and monitoring activities, either by a free search or by using predefined lists for biodiversity and socio-economic monitoring (Lotze-Campen et al. 2008). Studies focussing on the implementation of a BRIM are still scarce; Price et al. (2006) proposed a set of social indicators for mountain biosphere reserves, in order to (i) assessing the main trends and driving forces of human–nature interactions; (ii) contributing data for hypothesis testing and modelling; (iii) informing the evaluation of the contribution of biosphere reserves to sustainable development; and (iv) providing input to the policy cycle, more generally as part of the feedback process of policy analysis, and specifically, e.g. by providing mechanisms for early warning and information for the development of scenarios.

Figure 1 depicts the core idea of a BRIM, by representing the passage from observation of data coming from the natural and the social spheres to the phase of monitoring. The goal of the monitoring is to achieve sustainable management, in connection with existing objectives and current measures. For progressing in the monitoring phase, adequate indicators are needed. The necessity of sustainability indicators has been highlighted in previous studies (e.g. Banos-González et al. 2016), however further research efforts are required. This considered, in the framework of the development of a BRIM, the main tasks of scientific

research and analysis are indicator development. From this recognition takes shape the BRIMSEN project, since it provides a methodology for selecting a set of indicators that are inclusive (combining socio-cultural with environmental sphere), replicable and transferable to other biosphere reserves. Adequate indicators can be selected by making use of the ecosystem service perspective.

Figure 1: From observation to monitoring (Fischer-Kowalski, Erb and Singh 2004)

2.3 The concept of ecosystem services and of landscape services

It is nowadays globally recognized that biodiversity sustains human life by means of the so-called ecosystem services (e.g. Isbell et al. 2015). Ecosystem services are defined as ‘the conditions and processes through which natural ecosystems, and the species that make them up, sustain and fulfil human life’ (Daily 1997). They arise when a biophysical structure (e.g. vegetation cover) or function (e.g. slow passage of water) directly or indirectly contributes toward meeting a human need or demand. Such services (e.g. flood protection) generate benefits (e.g. serving health and safety) that contribute to overall well-being and can be valued by people (e.g. willingness to pay for protection) (Haines-Young et al. 2010; MEA 2005). According to the MEA (2005), an ecosystem function is an intrinsic ecosystem characteristic related to the set of conditions and processes whereby an ecosystem maintains its integrity (such as primary productivity, food chain, biogeochemical cycles). Ecosystem functions include such processes as decomposition, production, nutrient cycling, and fluxes of nutrients and energy. On the other hand, ecosystem services can be translated into the benefits people obtain from ecosystems. According to the Millennium Ecosystem Assessment (MEA 2005) they can be divided into provisioning services such as food and water; regulating services such as flood and disease control; cultural services such as spiritual, recreational, and cultural benefits; and supporting services such as nutrient cycling that maintain the conditions for life on Earth (MEA 2005).

The ecosystem service framework extends prior models by encouraging people to re-examine the links between ecosystems and human well-being in a more pragmatic way (Potschin et al. 2011a). Emphasizing ecological, social and economic dependence on natural

assets attracts public interest on biodiversity conservation (see Westman 1977; de Groot 1987; Odling-Smee 2005).

The ecosystem services approach is an established framework for the balanced evaluation of ecological, economic and social landscape resources. It promotes functional synergies (win-win situations) as well as trade-offs among various benefits resulting from ecosystem processes (Syrbe et al. 2012). Information, communication and knowledge are seen as crucial factors in promoting sustainable economic and social development as well as preservation of the environment at the local, national, regional and global levels. The concept of ecosystem service is considered promising for acting as a comprehensive communication tool.

Although the knowledge base on how to assess ecosystem services is improving, there are still substantial data gaps and further research is needed. The large number of projects concentrating on these issues e.g. TEEB (TEEB 2010), RUBICODE (Luck et al. 2009), BIOSERV (Hermann et al. 2014), TRANSECONET (Bianchin et al. 2012), ESMERALDA (e.g. Brown et al. 2018) is evidence of great interest in the progress of this research topic. At European level, the EU Biodiversity Strategy 2020 requires Member States to halt the loss of biodiversity and the degradation of ecosystem services in the European Union by 2020 (Target 2) (European Commission 2011). The actions designed to maintain and restore ecosystems and their services are: (i) improve knowledge of ecosystems and their services, (ii) set priorities to restore and (iii) promote the use of green infrastructure and finally (iv) to ensure no net loss of biodiversity and ecosystem services (Maes et al. 2016).

In BRIMSEN we addressed a variation of the concept of ecosystem services, i.e. landscape services. This means defining functions, services and benefits at landscape scale to integrate the concept into land management decisions (Bastian et al. 1999; de Groot et al. 2010; Willemen et al. 2010). Landscape services are the contributions of landscapes and landscape elements to human well-being (Bastian et al. 2014), and they include potentials, materials and processes of the nature (e.g. raw materials, biomass, biodiversity etc.) and services of cultural elements and constructions that come into being through human creation (e.g. buildings, settlements, infrastructure etc.) (Konkoly-Gyuró 2011; Hermann et al. 2011). The landscape services concept can be seen as a bridge between landscape and value. Since landscape sciences primarily focus on spatial pattern and scale, they offer valuable insights into the spatial distribution of human activities and their influence on important landscape processes and structures from which services are derived (Müller et al. 2008). Important reasons to consider landscape services include the prominent role of spatial aspects, the reference to spatial elements and to the landscape character, and the relevance of landscape services for spatial planning (Bastian et al. 2014). The pattern of multi-functional landscapes is the basis for interactions, synergies and conflicts between landscape elements (Willemen et al. 2012). Moreover, the provision of services does not always depend on the properties of an ecosystem patch, but rather on the spatial interaction among these patches (Termorshuizen et al. 2009). Last, as local people define their environment more as a “landscape” than as an “ecosystem” the term “landscape services” is preferred as a specification (rather than an alternative) of ecosystem services (Termorshuizen et al. 2009). Landscape services are supplied by landscape elements, i.e. the basic spatial entity of this study, in the sense of Forman & Godron (1986) and Forman (1995). Following the definition in this classic textbook of landscape ecology, landscape elements are “basic relatively homogenous spatial elements at the landscape scale” and may be “of natural or human origin, and thus apply to the spatial pattern of different ecosystems, community types, successional stages, or land uses”. In the central European tradition of applied landscape

ecology and conservation planning, the term “ecotope” or “biotope” has been often used (Kaule 1986), having a slightly different – more selective - meaning in the sense of habitats with a higher conservation value. Here, we prefer the more neutral term “landscape element” as any spatial element in the “open” agrarian part of the biosphere reserve, regardless its specific function as a potential habitat for endangered species.

To emphasise the political relevance of the ecosystem and landscape services concept, it is important to point out the additional value of ecosystem and landscape services produced by conserving an area with diverging utilisation claims. The importance of ecosystem and landscape services can be assessed in different ways. In the present project, we will address an assessment, which avoids monetary valuation, because the monetarisation of services would require modern economic valuation techniques, which would go beyond the scope of our project and beyond our fields of expertise. Furthermore, it was stated that economic valuation of services has reached its limits (e.g. Heal 2000; Farber et al. 2002; Wilson et al. 2002; Chee 2004; Hein et al. 2006). Moreover, an exclusive focus on the economic valuation of ecosystem outputs may narrow the debates around ecosystem services and deterring the development of the idea (Potschin et al. 2011b). It is important to realize that economic and especially monetary valuation will always capture only part of the “true” or total value (which should also include ecological and socio-cultural values) of an ecosystem or service (de Groot et al. 2010). Some researchers also suggest that a new approach to environmental policy is needed that goes beyond the traditional cost-benefit analysis (Ekins 2011).

Figure 2: Conceptual model showing relations of ecosystem functions, services and benefits (after Burkhard et al. 2014)

2.4 Service capacity and capacity matrix

Major challenges in all ecosystem and landscape service assessment efforts are the high complexity of the topic itself and the need for universal and rather easy-to-apply approaches (Crossman et al. 2013; Seppelt et al. 2012). In terms of conceptualization, there is still an ongoing debate on the interpretation of ecosystem and landscape services with respect to benefits, especially when applying the terms to landscape planning (e.g. de Groot et al. 2010).

To ease the terminological discussion, three concepts can be commonly distinguished in the ecosystem and landscape services framework: service capacity or potential, service flow and

service demand (Burkhard et al. 2014). Figure 2 shows the connections from the ecosystem functions, to the service supplies (potential and flow) till the fulfilment of human benefits. Service demand influences the ecological integrity of an ecosystem through land use, as the loop arrow clearly shows (Figure 2). A change in land use or management causes a change in service supply, not only for specific services but for the complete bundle of services provided by that ecosystem (de Groot et al. 2010).

BRIMSEN focuses on service capacity only, similarly to many of the currently available spatial ecosystem and landscape service studies (for instance Crossman et al. 2013; Martínez-Harms et al. 2012; Egoh et al. 2012). Service capacity can be here defined as the hypothetical maximum yield provided by a service (Burkhard et al. 2012; Burkhard et al. 2014). Assessing the actual capability of ecosystems to provide services for human well-being needs information about their current conditions which are induced by human activities (Burkhard et al. 2017). In this sense, nor the actual service capacity, neither the used stock of services is taken into account in BRIMSEN. Other authors distinguish between ecosystem properties, potentials and services (Bastian et al. 2012), implying that ecosystems provide a certain potential to supply services based on their functioning (van Oudenhoven et al. 2012).

A common method to assess the service capacity is the so-called capacity matrix. A capacity matrix links service providing units (defineable at different spatial scales) to ecosystem service supply capacities (Burkhard et al. 2009). In an assessment based on a capacity matrix, for each service providing unit a ranking proportional to the capacity for each service is assigned. The scores generally vary from 0 (no capacity) to 5 (very high capacity). Generally, expert evaluations are employed, in order to gain an overview and see trends for ecosystem service assessments (e.g. Burkhard et al. 2009; Scolozzi et al. 2012). In subsequent analyses, the expert evaluation values could successively be replaced by data from monitoring, measurements, computer-based modelling, targeted interviews or statistics, although these techniques imply a much longer data processing. By linking the 0–5 values to appropriate spatial data in GIS, estimates of ecosystem service supply can be mapped in spatially explicit units (Burkhard et al. 2009; Burkhard et al. 2012).

The method has successfully been applied to quantify and map ecosystem and landscape services in several case studies (e.g. Kandziora et al. 2013; Kaiser et al. 2013; Vihervaara et al. 2010; Vihervaara et al. 2012; Hermann et al. 2014; Hainz-Renetzeder et al. 2014; Stoll et al. 2015). It has also inspired the development of ecosystem service mapping studies (e.g. Clerici et al. 2014; Baral et al. 2013).

3 Research questions

In BRIMSEN we focus on the development of a monitoring tool for biosphere reserves which can be used for the systematic evaluation of goal attainments. We provide a toolset of indicators as a basis for sustainable development focusing on nature conservation, some economic activities, and landscape related recreational activities within the Wienerwald Biosphere Reserve. Our indicators are based on the concept of landscape services, which can be seen as policy-relevant representations to identify gaps and to communicate trends in a sustainable landscape development (Layke et al. 2012). In order to achieve our goal, we set out to answer the following research questions:

- Which landscape elements are suitable for representing the landscape services of the open land taking into consideration the availability of spatial data?
- How to simultaneously assess and spatially represent ecological and socio-cultural landscape services?
- How can the landscape services assessment contribute to a replicable monitoring of the Biosphere Reserve sustainability performance?
- How do the landscape services spatially vary and interact within the Biosphere Reserve?
- Which landscape services can be interpreted as indicators for a BRIM?

4 Material and Methods

4.1 Project area description

The Wienerwald Biosphere Reserve (designated as Biosphere Reserve by UNESCO in 2005) encompasses an area of 105.645 hectares within the Austrian provinces Lower Austria and Vienna (Figure 3). Influenced by the easternmost part of the Alps, it is characterized by a hilly terrain (sea-level from 160m up to 893m). Geologically the Biosphere Reserve can be divided into the northern part consisting mostly of flysch rock ("Sandstein-Wienerwald") and the southern part, primarily limestone ("Kalk-Wienerwald") that appears near the so-called "Thermenlinie". As a stepping stone within the biotope network the Wienerwald is of superregional importance (Loiskandl 2005; Reimoser et al. 2008; Reimoser et al. 2012)

The open land areas of the Biosphere Reserve consist of 23 different types of meadows; most common is the oat-grass meadow ("Glatthaferwiese"). Together, the meadows and pastures have a share of 12% of the Wienerwald Biosphere Reserve area. The climatic, geological and biogeographical border location, special habitats as well as close interaction between areas of forest and open land (like meadows, pastures, cultivated fields, vineyards etc., including elements like single trees, orchards, hedge and tree rows) allow for a diverse mosaic display of the Wienerwald landscape, increasing the ecological and nature conservation value of the Biosphere Reserve (Reimoser et al. 2008; Reimoser et al. 2012; Drozdowski et al. 2018). More than 60% of the Biosphere Reserve area is covered by forest (Loiskandl 2005; Brenner et al. 2015). Based on climatic and geological characteristics such as sub-continental cold winters, dry and hot summers and associated soil qualities, the forests are mainly composed of oak and hornbeam, but about 33 different types of forests occur in the area (Brenner et al. 2015; Reimoser et al. 2008; Reimoser et al. 2012). In the Wienerwald Biosphere Reserve nature conservation, economic and social development as well as preservation of cultural values play an important role. It includes 27 core areas with particularly valuable forests, which equate 5% of the Biosphere Reserve area. The buffer areas, 19% of the Biosphere Reserve surface, contain valuable cultural landscape of meadows and grassland (Biosphärenpark Wienerwald Management GmbH n.d.)

The Wienerwald Biosphere Reserve intersects with seven municipal districts of Vienna as well as 51 municipalities in lower Austria: whereas in three municipalities in lower Austria live more than 20.000 inhabitants each, 60% of the municipalities have less than 5.000 inhabitants; altogether the Biosphere Reserve region hosts 815.000 inhabitants (Biosphärenpark Wienerwald Management GmbH n.d.). Beside the agricultural and forestry-used area, there is pressure from settlements, infrastructure and recreation. Because of the settlement areas, the proximity to Vienna and the characteristic landscape, the Biosphere Reserve is strongly used by recreationists. The area offers possibilities for various recreational activities and sports and is further enriched by a broad range of tourist attractions of cultural value (Loiskandl 2005; Reimoser et al. 2008; Reimoser et al. 2012).

For the BRIMSEN project the focus of the study area was on the open land areas of the Wienerwald Biosphere Reserve exclusively (Figure 3), which corresponds to an area of 27.831,5 hectares within the Biosphere Reserve. Due to its high potential for a wide spectrum of services, the forest was included by integrating the forest edges into the analysis. Furthermore, since the focus is more on the pull factors of recreation users within

the open land, rather than push factors, settlements were excluded from the analysis. Therefore, the potential of the service “habitation” is approached through the indicator “isolated buildings”. A difference in age and accuracy of the geodata underlying the project analyses required a quality-based selection of data within the south eastern region along the border of the Wienerwald Biosphere Reserve. On these grounds, part of the south eastern region of the Biosphere Reserve was excluded, fact that shall be considered in the valuation of the regional assessment of the landscape services.

Source: NUTS - Eurostat, 2018; Staudinger et al. (2014). Report - Methodology of the project open land survey BR Wienerwald 2011 (part 1) and 2012-2013 (part 2)

Figure 3: Study area: the open land of the Wienerwald Biosphere Reserve.

4.2 Data sources

The project main data source is the open land data of the Wienerwald Biosphere Reserve, originating from a detailed mapping of open land habitat types performed in the field mapping project “Offenlanderhebung Biosphärenpark Wienerwald” (Staudinger et al. 2014) for the open land of the whole Biosphere Reserve in Lower Austria and Vienna. The data was provided by the management of the Biosphere Reserve and is here further referred to as “open land data”. The geometries are based both on cadastral maps with high accuracy and remote sensing with the accuracy ranging between 10-30m (Schrantz 2018, pers. comm.). Differences in the age and accuracy of the open land data in the southeast of the Biosphere Reserve region, along the so-called “Thermenlinie”, led to a bias within the calculation of the results. Therefore, we decided to exclude the data based on the selection query: “open land data AND within Lower Austria AND no mapping date”. The exclusion was performed in a GIS environment. Possible inaccuracies within the open land data mapping arising from changes in the cultivated landscape over time e.g. the current condition of fallow land, were not revised. No further data verification of the open land data was carried out.

We relied on the open land data as the project prime data source and included data from other data sources solely when additional information was needed to address a service more fully or if it provided more detailed information on transformed landscape elements than the natural element-focussed open land data. All additional data was customised to the spatial projection of the open land (EPSG: 31259) using ArcGIS 10.5.1 (ESRI, Redlands).

The additional spatial data (see section 4.4.2) was downloaded from OpenStreetMap provided by Geofabrik (Geofabrik GmbH et al. 2019). This data was used to extract information on polylines as roads or railways and additional points as religious buildings and other religious elements. The recreational and educational trails (polylines) originate both from the detailed data of the open street map as well as *.gpx data available on the touristic

website of the Wienerwald Tourismus GmbH (Wienerwald Tourismus GmbH 2018) for the mountain bike routes and educational trails. The data sources of the additional spatial data are listed in Table A 2.

Furthermore, we received the outer border of the Wienerwald Biosphere Reserve as a Shapefile from the Wienerwald Biosphere Reserve Management staff.

4.3 Definition of landscape services addressed in the project

In BRIMSEN we defined the landscape service classification addressed based on the definitions developed by de Groot (2002; 2006), thereby following the projects BIOSERV (Hermann et al. 2014) and CCR (Hainz-Renetzeder et al. 2014) in the Austrian-Hungarian Biosphere Reserve Neusiedler See. The concept of landscape services supports the interdisciplinary ecological and socio-cultural approach of BRIMSEN and it refers strongly to the cultural landscape and landscape elements incorporating natural and cultural aspects of the Wienerwald Biosphere Reserve (Bastian et al. 2014). The concept is also transferable to the Biosphere Reserves planning and management processes. For more details on the landscape services see section 0. The classification proposed by de Groot (2002; 2006) includes 25 landscape services distinguished into five service categories: **Regulation, Habitat, Provisioning, Carrier and Information** (Table A 1).

We adapted the list of landscape services definitions to the properties of the study area:

- Raw materials: the Raw material service includes the provision of sand and gravel e.g. by periodic brooks.
- Cultivation: the Cultivation service refers only to the provision of substrate for the cultivation of food or fodder, not for ornamental cultivation (e.g. gardening). Furthermore, in comparison to the service Food, it refers to cultivated fields (producing crops and fodder), vineyards and orchards as well as managed meadows (fodder production)
- Food: the service Food refers to all food produced for human use (wild and cultivated) and includes all animal related farming (aquaculture, cattle etc.).
- Genetic resources: in BRIMSEN we extend the Genetic resources to forestry and agriculture.
- Waste disposal: in our definition the service includes the landscape elements providing wastewater disposal.
- Transportation: Potential transportation on waterways was not included in the evaluation.
- Tourism facilities: as a carrier service, transformed (man-made) landscape is addressed exclusively.
- Recreation: mainly referring to natural landscape elements used for recreational purpose. Furthermore, following de Groot's inclusion of eco-tourism within the Recreation service, recreation infrastructural elements are also included in the service (de Groot 2006).

Although we addressed the classification proposed by de Groot (2002; 2006), many other classifications of ecosystem services exist. The most well-known classification was defined in the framework of MEA (2005). Other relevant classifications are those proposed by TEEB (2010), Müller and Burkhard (2007), and CICES (Haines-Young et al. 2018). Müller and Burkhard (2007) integrate the concept of ecological integrity as pre-condition for the supply of regulating, provisioning and cultural ecosystem services. The CICES (Common International Classification of Ecosystem Services) was first released in 2013; it offers a

structure that links with the framework of the UN System of Environmental-Economic Accounts (United Nations 2003), and it has become nowadays the most standardized classification at EU level (Maes et al. 2016).

In order to link the BRIMSEN conceptual approach to the EU standardized classification, we compared the BRIMSEN landscape services definitions with the ecosystem service classification CICES V5.1 (Haines-Young et al. 2018). As an element for comparison, we decided to adopt the “Simple descriptor” available in CICES V5.1, since such descriptor provides an unambiguous definition for most of the services. For the comparison we adopted a multiple correspondence approach. The selective, multiple correspondence approach (Figure 4) allows for more than one CICES ecosystem service to be linked to different landscape services without the need of assigning a landscape service to every CICES ecosystem service.

Figure 4: Scheme of the approach for comparing the CICES and BRIMSEN classifications.

Table A 1 reports the definitions used in BRIMSEN, the corresponding definitions according to CICES (Haines-Young et al. 2018) and examples of goods and benefits. The definition of the landscape services serves as a basis for the development and valuation of the landscape element classes considered in BRIMSEN. The comparison of the landscape services with CICES ecosystem services indicated that the landscape carrier services are underrepresented in CICES, emphasising the natural environment and giving less importance to the transformed landscape elements and to the socio-cultural aspects. On the other hand, CICES proved to be more detailed in distinguishing the regulating and provisioning ecosystem services, like for instance Disturbance prevention, for which four simple descriptors could be identified, and Food, for which seven simple descriptors could be identified (Table A 1).

4.4 Landscape elements classification

In BRIMSEN, we refer to landscape elements as our smallest entity, including both natural and human modified landscape components (Forman et al. 1986; Forman 1995). To assess the landscape elements, they were assigned to classes.

4.4.1 Open land habitats

As mentioned in section 4.2, the spatial source of the open land habitats is the survey and detailed mapping of the open land habitat types within the biosphere reserve finalized in 2014 (Staudinger et al. 2014). We reclassified and aggregated the open land habitats, so to obtain a list of open land classes, representative for both ecological and socio-cultural landscape services.

The reclassification of open land habitat types into open land classes was achieved in three steps (Figure 5). First, in an ecological approach, the open land types were aggregated into classes, based on management level, nutrient level, moisture level, and ecosystem function. In a second step, taking socio-cultural aspects like recreational function or aesthetic aspects into account, the classes were re-composed. Finally, the two approaches evolved into one joint reclassification, consisting of 62 different open land classes e.g. straight semi natural streams, reed beds, species-poor field margins, tree rows and single trees (see Table A 3). Homogeneous open land groups were then formed by aggregating more classes. Each group was identified through the first two letters of the unique code assigned to each open land class (Table A 3).

The hierarchy of the reclassification, exemplified in Figure 5, follows the structure: landscape elements → open land types (referring to the original habitat list used in Staudinger et al. (2014) → open land classes (the combined types) → open land groups (overarching group of open land classes).

Figure 5: Example of open land classification (OL= open land).

The study area exclusively focusses the open land areas of the biosphere reserve on a very detailed level:

- Due to its high potential for a wide spectrum of services, the forest was included by integrating the forest edges into the analysis.
- The focus on the pull factors of recreation users within the open land, rather than push factors, allows for an exclusion of the settlements. Therefore, the potential of the service Habitation is approached with the open land class “isolated buildings”.
- Due to discrepancies in the mapping quality of the data, a selection of the open land data of the south eastern area of the Wienerwald Biosphere Reserve was necessary. Therefore, not all the open land elements in the south eastern area are included in the calculations (see section 4.2).

4.4.2 *Additional spatial indicators*

Lying spatially within the above-defined natural landscape elements of the open land, a range of additional transformed landscape elements (roads, trails, castles, churches etc.) was included. To allow a more detailed socio-cultural approach these landscape elements were selected to add information and insight mainly to the **Carrier** and **Information** services of the Biosphere Reserve, since they were not adequately represented by the open land habitats. Similarly to the approach employed for the open land, the elements selected as additional spatial indicators were classified, based on their similarity and on their service provision capacity, into 18 additional spatial indicators e.g. public transport, religious buildings, museums, recreational infrastructure (see Table A 4).

The reclassification of the open land habitats into 62 open land classes and the selection and classification of additional spatial indicators into 18 classes led to a joint collection of 80 indicators classes (including altogether 58.202 landscape elements), which, in a next step, were inserted in a capacity matrix for the landscape service capacity assessment.

4.5 Capacity matrix: assessing service capacities

Having set a landscape service categorisation and having developed the indicator classification, we assessed the capacity of the 25 landscape services for each of the 80 indicators classes. Following the works of Burkhard et al. (2009; 2012; 2014), Hermann et al. (2014), and Hainz-Renetzder et al. (2014), we applied the following scoring scheme, in a linear scale:

- 1: element presence, but no service capacity
- 2: low service capacity
- 3: moderate service capacity
- 4: high service capacity
- 5: very high service capacity

The 25 landscape services were split and assigned to the two teams of expertise (ecological and socio-cultural) involved in the project. First, both teams of experts assigned values for the according services, producing a prefilled matrix. Then, through a dedicated workshop all experts contributed to the assessment of the values, by discussing the scores of the services in their field of expertise and by providing comments and suggestions regarding services of the other discipline.

Helpful within the workshop was the preparation of an approachable layout of the matrix in the software Excel (see Table 1). To ensure transparency, the matrix was projected on a screen, clearly visible for all people involved. We approached each class (e.g. straight, seminatural streams) individually, discussing and comparing the values vertically (to other classes) and horizontally (to other landscape services). When a consensus was necessary, we adjusted the scores directly on the screen, again aiming for high transparency. Provision of the definitions of landscape services and information on the open land types within the classes was necessary. The possibility to examine the location of the classes within the geodata and an orthophoto were helpful to agree on the scores. The orthophoto was applied for inexplicit values only, due to the time-consuming procedure.

Altogether, 2000 scores were given within the BRIMSEN joint capacity matrix of 62 open land and 18 additional spatial indicators classes addressed within 25 landscape services.

Table 1: Extract of BRIMSEN capacity matrix.

Open Land Class	Science and education	4	4
	Spiritual and historic information	4	5
	Cultural and artistic information	2	3
	Recreation	2	4
	Aesthetic information	3	5
	Information services		
	Tourism facilities	1	1
	Transportation	1	1
	Energy conversion	1	1
	Habitation	1	1
	Waste disposal	3	3
	Cultivation	1	1
	Carrier services		
	Medicinal resources	1	1
	Genetic resources	2	2
	Raw materials	2	2
	Food	3	3
	Provisioning services		
	Nursery	5	5
	Refugium	4	4
Habitat services			
Pollination	2	2	
Nutrient regulation	5	4	
Soil formation	2	1	
Soil retention	1	1	
Water supply	5	5	
Water regulation	5	2	
Disturbance prevention	5	4	
Local climate regulation	5	4	
Regulation services			

The BRIMSEN capacity matrix scoring gives no absolute values since the scores refer to a local approach of the Wienerwald Biosphere Reserve. The comparison of indicators inside the area influences the value assigned. For instance, compared to dry grassland, other grassland has higher water supply; this means that peaks can be developed within the matrix. As described in section 2.4, the evaluation focusses on the potential capacity of the open land. Agreement on definitions to describe each landscape service helped to avoid issues of double counting, which refer rather to the service, than the indicator (Bianchin et al. 2012). In case of similar aspects being valued within two services, values were split

between the two (e.g. recreational infrastructure is valid both for landscape services Tourism facilities and Recreation).

The collection of expert knowledge within our capacity matrix represents an important research output and essentially contributes to the creation of the capacity maps of the Wienerwald Biosphere Reserve, described in the following sections.

4.6 Spatial data preparation

Simultaneously to the scoring of landscape service capacity, we prepared the spatial data so to later integrate the capacity matrix scores in the mapping process.

The forest-related classes of the open land dataset were transformed into polylines, since the forest was addressed as forest edge exclusively.

With reference to the accuracy of the open land data, a buffer of 10m was applied to the data to secure the inclusion of all point and line features within or in direct proximity of the open land. By clipping the buffered data to the Wienerwald Biosphere Reserve's outline, we defined this as the outermost border of the project study area. The geodata processing was done using ArcGIS 10.5.1 (ESRI, Redlands).

Due to the "layman" and inclusive nature of the OpenStreetMap data (OpenStreetMap Austria 2019), a deletion and ranking of the point data with congruent location within the study area was indispensable. Referring to the precise data, expert decisions towards a double function of the element or towards its deletion were made. Hence part of the double counted data was dismissed due to (i) redundancy (e.g. graveyards data was adopted from the open land class; "convenience stores" or "wind mill" were not included in the dataset); (ii) ambiguity (the vague meaning of "attractions" was recessive). In case of "church" and "wayside cross" we decided in favour of the smaller element. We decided a picnic site to be an inclusive element for a bench or a waste basket. Other OpenStreetMap elements provided a double function; therefore, both points were included in the further calculations for their double potential (e.g. "guesthouse" and "restaurant", "bench" and "viewpoint", "restaurant" and "viewpoint", "church" and "artwork", "hiking" and "cycling trail").

4.7 Transformation of spatial data

Due to the heterogeneity of data sources and data sets (spatial resolution and temporal variance) and the ambiguity in size for some of the additional spatial indicators (e.g. streets, playgrounds, churches), we developed a methodology that would overlook both the size and the differences in data sources. The approach was based on two concepts: location of elements and the capacity scores retrieved directly from the capacity matrix. For the location of the landscape elements all the data types (line, polygon, and point) were transformed into point data. Figure 6 displays the workflow of the calculation. First, we prepared the spatial data to be transformed into points, and then we applied an innovative tool for mapping the landscape service capacity, which we named **INDEC** (analysis based on **IN**tensity and **DE**nsity of service **C**apacity). Each passage of the workflow is described in the following paragraphs.

Figure 6: Workflow of the INDEC calculation, with data preparation and INDEC application.

In order to transform the polylines into points, we had to compute the point distance. In fact, no references for setting point distance could be found. For the line features the equivalent of the sum of all the lines within the area (5.027.130m) was equally distributed into an abstract quadrad of the research area size. Within this generated grid, we doubled the value of the side lengths of one grid cell (223m), thereby prohibiting double counting mistakes leading to a misrepresentation of the line features. This value represents the ideal and equal distribution of points for the line features, similar to the method used for transforming polygons into points (Figure 7).

Calculation of the distance of points along line features

d_{points} = distance of points along polylines
 a_{research} = research area
 l_{total} = total length of all polyline landscape elements
 $d_{\text{points}} = 2 * (2 * a_{\text{research}} / (l_{\text{total}} - 2 * a_{\text{research}}^{1/2}))$

Figure 7: Scheme for point distance calculation

Regarding the polygons, we transformed them into a point coinciding with the focal points of the polygons. Subsequently, the points generated from different geometrics (point, polyline, and polygon) and sources were merged into one “point feature class”.

Based on the assumption that an ideal landscape services map would show an equally distributed maximum capacity of the service, we computed the equally distributed area of each point data (4.871m²) from the total study area (278.315.000m²) and the number of landscape elements (58.202).

Calculation of the maximum buffer size of LE

bs_{max} = maximum buffer size

$a_{research}$ = research area

$n_{LE-total}$ = total number of all landscape elements (points)

$$bs_{max} = a_{research} / n_{LE-total}$$

In this theoretical optimum, each landscape element has the same maximum capacity to provide a landscape service and the elements are equally distributed across space. The calculated areas are perceived as circles (see Figure 8). The circles represent each landscape element with an equal range in all directions from one centre point, based on the assumption that the capacity of one landscape element potentially spreads equally in each direction. Therefore, the capacity of each point is represented by an outer buffer, whose radius is linked to the scores from the capacity matrix. The buffers describe the potential intensity of the capacity. We computed the radius of the average circular area and adopted it as size of the maximum capacity buffer (equal to 5 in the capacity matrix). At the maximum service capacity, the buffers (radius = 39m) partially overlap (Figure 8). In this way the connectivity between the landscape elements is guaranteed.

Figure 8: Graph exemplifying the theoretical assumption for defining the maximum buffer size (39m) of the landscape service capacity of a landscape element.

For defining the intensity related to each point, we calculated the size of the buffer of the remaining capacity matrix scores (2-4) related to the maximum area (4.871m²), and not to the radius, thus again assuring the connectivity of the capacity buffers. E.g. 50% capacity (value 3) of maximum capacity (value 5) is equal to 50% (2.435,5m²) area of maximum area (4.871m²).

- 1: presence, no capacity. Data with score 1 was excluded from the calculation.
- 2: low capacity. It equals 25% of the maximum capacity.
- 3: moderate capacity. It equals 50% of the maximum capacity.
- 4: high capacity. It equals 75% of the maximum capacity.
- 5: very high capacity. It equals 100% of the maximum capacity.

Calculation of the relation between buffer size and capacity value

b_{r-5} = buffer radius of landscape elements with capacity 5

bs_{max} = maximum buffer size

$b_{r-5} = bs_{max}$

b_{r-x} = buffer radius of landscape elements with capacity x (e.g. 2-4)

b_{r-1} = buffer radius of landscape elements with capacity 1

x = capacity

$$b_{r-x} = b_{r-5} * ((x-1)/4)^{1/2}$$

$$b_{r-1} = 0$$

The spatial data preparation provided two outcomes: (i) the transformation of each landscape element into point data; and (ii) definition of the buffer sizes according to the scores given the capacity matrix.

4.8 INDEC application

After the preparation of the spatial data and the transformation to point data for all landscape elements, we applied the newly created tool named INDEC.

First, we linked the landscape elements to the scores provided for each service in the capacity matrix. For the polygons we included an area weighting factor, which refers to the biggest (e.g. cultivated field ca. 400.000m²) and smallest (e.g. single tree, a few m²) landscape element of the open land. The relative part of the maximum area became 1, and the relative part of the minimum area became 0. All the remaining elements received relative values between 0 and 1. The relative part allows for a ranking within the elements, according to the relative size (referring to the largest polygon).

Calculation of the relative part of each LE size

rp_{LE-x} = relative part of LE x (one open land LE e.g. meadow)

a_{LE-x} = area of LE x

a_{LE-min} = area of the smallest LE

a_{LE-max} = area of the biggest LE

$$rp_{LE-x} = (a_{LE-x} - a_{LE-min}) / (a_{LE-max} - a_{LE-min})$$

We transformed the relative part of the landscape elements to a weighting factor, depending on three defined maximum weighting factors, by linear correlation.

As depicted in Figure 9, depending on the relative part (0-1, 1 equal to ranking position 1) and on the defined maximum weighting factor (wf) each landscape element receives an adequate wf. For example: in case of relative part "0,5" and maximum weighting factor "5", the transformed weighting factor is "3".

Calculation for the transformation of the relative part to weighting factor

wf_{LE-x} = weighting factor for landscape element x

wf_{max} = maximum weighting factor

rp_{LE-x} = relative part of landscape element x

$$wf_{LE-x} = (wf_{max} - 1) * rp_{LE-x} + 1$$

Figure 9: Calculation of weighting factor based on relative part.

For the line elements the same relative size calculation was done referring to an expert-set distribution factor (see section 4.9.4) based on the width of the line elements.

In a geoprocessing step, we buffered the elements according to the values computed by multiplying the buffer value (from the capacity matrix score) with the weighting factor of the line and polygon landscape elements. We merged the overlapping buffers by “spatial dissolve”, obtaining clusters (dissolved shapes) representing the cumulative intensity of service capacity. Subsequently we counted the number of landscape elements within each cluster by “spatial join”, using this number as an indicator of the *density* (**INDEC**) of service capacity.

Finally, we combined both *intensity* (the size of the cluster) and *density* (the number of elements within the cluster), creating our final **INDEC**. The **INDEC** itself refers to no unit and originates by 50% from the relative size and 50% from the relative amount of the landscape elements. All landscape service capacities are calculated through the same **INDEC** procedure, although for each landscape service only the landscape elements with service capacity (capacity matrix values 2-5) were included in the processing.

For the index of the intensity ($I_{IN-cl-x}$) of a cluster, the cluster area (A_{cl-x}) is divided by the normalized cluster area ($A_{cl-norm}$) which corresponds with the optimum area per landscape element ($A_{oALE} = bs_{max} = 4.871m^2$). This value refers to the value of the equally distributed area that is used to calculate the maximum buffer size and now allows for a normalisation as reference of the area size. In case the cluster size exceeds the optimum area per landscape element, it will produce a value higher than 1. In case of smaller clusters, a value lower than 1 will be produced. For clusters equal in size to the optimum area per landscape elements, the value 1 is assigned.

For the index of the density ($I_{DE-cl-x}$) the number of elements within the cluster ($n_{LE-cl-x}$) is divided by the number of elements per optimum area of landscape elements ($n_{LE-per-oaLE}$)

equalling 1. The sum of the index of intensity ($I_{IN-cl-x}$) and the index of density ($I_{DE-cl-x}$) is the combined INDEC ($I_{INDEC-cl-x}$).

Calculation of the final value based on intensity and density of capacity (INDEC)

$A_{cl-norm} = A_{oaLE}$ = optimum area per LE (bs_{max})

$n_{LE-per-oaLE} = 1$ = number of LE per optimum area of LE

A_{cl-x} = size of cluster x

$I_{IN-cl-x}$ = Index of intensity of cluster x

$I_{DE-cl-x}$ = Index of density of cluster x

$I_{IN-cl-x} = A_{cl-x} / A_{cl-norm}$

$I_{DE-cl-x} = n_{LE-cl-x} / n_{LE-per-oaLE}$

$I_{INDEC-cl-x} = I_{IN-cl-x} + I_{DE-cl-x}$

4.9 Refinements to the methodology

After creating the preliminary capacity maps, we refined the applied method to obtain higher representativeness of each service capacity.

4.9.1 Exclusion of landscape services

Data availability was a limiting aspect within the project. By using different data sources (see chapter 4.2) we expanded our database to provide some indicators for all the services. However, after visualizing the preliminary capacity maps, three services had to be excluded from our list of landscape services, due to incomplete representativeness. The three services were the following:

- Habitation: Exclusion of the service due to differences in the accuracy of the data. The mapping of the habitation related data collected in Staundinger et al. (2014) did not match the detailed precision given to habitat types of the natural open land. Other data sources (e.g. open street map) were not compatible with the open land data. Therefore, the service was dismissed.
- Waste disposal: Lack of data led to the exclusion of the service from the assessment.
- Energy conversion: The available geodata for the service was limited to the existing hydro power plants (provided by Institute of Hydrobiology and Aquatic Ecosystem Management) within the Wienerwald Biosphere Reserve, which was not sufficient information to assess the energy conversion service at regional scale. Other types of energy conversion units (e.g. wind turbines) do currently not exist in the Biosphere Reserve.

4.9.2 Validation of plausibility of the landscape services capacity maps

The plausibility of landscape services capacity maps was validated with expert knowledge of the area during an internal workshop. By supplying preliminary maps and devices with the data displayed in Google Earth Pro (Google Earth Pro n.d.), two groups of experts from the team involved in BRIMSEN were tasked to provide information on the areas of high and low capacity and their location within the map of each landscape services. The discussion following the individual tasks revealed three main aspects for refinement:

- Due to the spatially scattered nature of the study area, the weighting factor was further adjusted (preliminary maps received weighting factor 2) in order to display high visibility of the service capacities on the regional level.
- Some line elements (forest edges) appeared overrepresented, creating a bias in the results, and therefore received a separate weighting.
- The scoring of the capacity matrix was revised to highlight some open land and additional spatial indicators classes with respect to others.

Considering the outcome of the plausibility task, we performed the amendments described in the following sections.

4.9.3 *Weighing factor adjustment*

By testing and evaluating the effects of different weighting factors on the capacity map results of a diverging selection of services (landscape services Water regulation, Refugium, Cultivation, Transportation, Recreation, Tourism) referring to diverse range of landscape elements, we approached the decision on the final weighting factor. On account of a plausible and balanced display of the landscape services addressed, the final BRIMSEN weighting factor was set to 6 for polygons and 3 for polylines.

4.9.4 *Treatment of polylines*

We approached the issue of strong representation of polyline elements (e.g. forest edges) compared to other line elements (e.g. primary roads) and to polygon elements of the open land (e.g. cultivated fields) on two different levels:

First, we provided a distribution factor referring to the hierarchy based on the width of the elements. Merely underrepresented elements received an increased value (2 or 3); others remain unchanged (value 1).

Table 2: Width distribution factor applied to polyline landscape elements (LE).

Polyline LE	Width distribution factor
Public transport	1
Roads primary	3
Roads secondary	2
Roads tertiary	1
Roads residential	1
Recreational trails	1
Educational trails	1
Forest edge	1

Second, we decided to assign the polyline landscape elements a different maximum weighting factor (3) with respect to the polygons (6), as mentioned above. In this way, we were able to regulate the representation of lines within our method, based on their nature of being represented by many points, in comparison with the polygons, which can only be represented by one point (i.e. their focal point).

4.9.5 *Validation of the capacity matrix*

The preliminary results of some landscape services required adjustments to the scores of the capacity matrix. Therefore, each group of experts revised the scores within their field of expertise and contributed again to a joint capacity matrix.

Changes affected 8% of the scores within matrix (2.000 scores in total), with 5% being upgraded to a higher score, and 3% being downgraded. The services Medicinal Resources, Disturbance prevention and Nutrient regulation faced the greatest changes with adjustments of 19% and 24% of the scores, respectively.

With these refinements we finalized our method and produced the final version of the service capacity maps, presented in the section 5.2.

4.10 Selection process of landscape services for BRIM

The selection of landscape services for a BRIM followed three stepwise criteria (represented in Figure 10):

Representativeness: The landscape services for BRIM shall be represented by a high quantity of landscape elements with high capacity within Wienerwald Biosphere Reserve. For each of the 22 landscape services we computed a cumulative score from the quantity of landscape element classes with capacity (values 2 to 5) and the amount of classes with high capacity (values 4 and 5) addressed in the capacity matrix (Figure 10, left column). The 16 highly ranked services (scores 4-6) provided the basis for the selection of the next criteria (Figure 10, left column, red box).

Communication: To assure the BRIM understandability for the management of the Biosphere Reserve and respective stakeholders, we further selected the landscape services conveying a clear message without ambiguity. We short listed the 10 landscape services (Figure 10, middle column) Local climate and air regulation, Refugium, Nursery, Soil retention, Water supply, Recreation, Aesthetic information, Science and education, Spiritual and historic information, Food. These 10 landscape services served as a pre-selection for the next criteria.

Relevance: The final selection comprises landscape services with high relevance for the management by providing information regarding the different spheres (Anthroposphere, Atmosphere, Lithosphere, Hydrosphere, and Biosphere) and thereby addressing the environment of the Wienerwald Biosphere Reserve as a whole. Emphasising landscape services with overarching character, but also considering a balance between ecological and socio-cultural aspects, we selected the 8 landscape services (Figure 10, right column) Local climate, Refugium, Soil retention, Water supply, Recreation, Aesthetic information, Science and education and Food.

Figure 10: Process of landscape services selection for a BRIM.

5 Results

5.1 Capacity matrix

The capacity matrix consists of landscape elements classified into 80 rows of classes, of which 62 are open land classes and 18 are additional spatial indicators. Out of the 25 landscape services, 22 landscape services were selected and arranged into 5 service groups. The matrix is expert-based and was developed by two groups of experts, addressing both the ecological and socio-cultural landscape services within the Wienerwald Biosphere Reserve.

The evaluation focusses on the potential capacity of the class for the landscape services.

The additional spatial indicators provide capacity for the landscape services categories **Information** services, **Carrier** services and **Habitat** services. The open land classes are addressed by all five service categories, but only marginally by the **Carrier** services.

In the evaluation of some classes (e.g. trails, recreational infrastructure) the score within the matrix was split, due to the importance of the elements for both services.

In the following pages the BRIMSEN capacity matrix will be presented group-wise, following the chronological order of the open land and additional spatial indicators classes.

Table 3: Capacity matrix of the open land (OL) classes 1-11

OL Code	Open Land Class	Regulation services	RS1	RS2	RS3	RS4	RS5	RS6	RS7	RS8	Habitat services	HS1	HS2	Provision services	PS1	PS2	PS3	PS4	Carrier services	CS1	CS5	CS6	Information services	IS1	IS2	IS3	IS4	IS5
WT1	Straight, seminatural streams		5	5	5	5	1	2	5	2		4	5		3	2	2	1		1	1	1		3	2	2	4	4
WT2	Meandering, seminatural streams		5	5	5	5	1	2	5	2		4	5		3	2	2	1		1	1	1		5	4	3	4	5
WT3	Channel/regulated streams		5	4	2	5	1	1	4	1		3	3		2	1	2	1		1	1	1		3	2	2	2	3
WT4	Periodic/small brooks		4	5	4	4	1	2	4	1		4	4		2	2	1	1		1	1	1		3	3	2	2	3
WT5	Seminatural, nutrient-poor lakes and pools		5	4	4	5	1	1	4	2		4	5		3	1	2	1		1	1	1		4	3	3	3	4
WT6	Nutrient-rich pools		5	4	4	5	1	1	3	2		3	5		4	1	2	1		1	1	1		2	3	2	2	3
WT7	Artificial standing water connected to groundwater		5	4	4	5	1	1	4	2		3	4		3	1	2	1		1	1	1		2	4	2	2	3
WT8	Artificial pools disconnected from groundwater		5	3	2	3	1	1	3	2		3	4		3	1	2	1		1	1	1		2	3	2	2	3
WT9	Springs, marshes, swamps		4	4	5	5	2	4	5	3		5	4		2	1	1	3		1	1	1		3	1	1	2	3
WT10	Nutrient-poor fens		4	3	5	5	3	4	5	4		5	4		2	2	1	4		1	1	1		2	1	1	1	1
WT11	Reed beds		5	4	5	4	4	5	5	3		5	5		2	3	2	2		1	1	1		4	1	1	1	2

Table 3 includes the first 11 classes for which the landscape services assessment was carried out. These classes are all water-related, as they include streams, both semi-natural and channelled, pools, and wetlands (springs, marshes, swamps, nutrient-poor fens, and reed beds).

All these classes have high or very high capacity for Local climate regulation, Disturbance prevention, Water regulation, Water supply, and Nutrient regulation. The other **Regulating services**, such as Pollination and Soil formation and Soil retention, showed medium or low capacity for these classes, with wetlands excluded.

Artificial water-related classes, as artificial standing water disconnected to groundwater and channel/regulated streams show lower water regulation capacity than the more natural classes. Nutrient-poor fens have the highest Pollination capacity in the water-related classes. Shifting to **Habitat services**, all water-related classes have medium to very high capacity for both Refugium and Nursery. Again, more artificial classes showed lower scores compared to the semi-natural classes.

Provisioning services are under-represented in water-related classes, with the exception of nutrient-rich pools, which have high Food capacity, and nutrient-poor fens, which have high Medicinal resources capacity. Food received more medium scores than the other **Provisioning** services, which have no or low capacity.

The water-related classes have no capacity for the landscape services of the **Carrier** service category.

Within the group of **Information** services, most water-related classes are valued with low or medium capacity (scores 2-3), except for the seminatural streams which have high capacity (score 4-5) for landscape services Spiritual and historic information, due to the historic influence of water courses to the landscape development and landscape services Science and education. The importance of natural and seminatural classes and their potential species richness for scientific work and nature education is also visible in the high score of the seminatural pools and lakes. The meandering, seminatural streams, the seminatural lakes and the reed bed also have high values (scores 4-5) for the landscape services Aesthetic information. The potential accessibility of the riverbanks of a meandering stream justifies the high capacity (score 4) for the landscape service Recreation.

Table 4: Capacity matrix of the open land (OL) classes 12-24

OL Code	Open Land Class	Regulation services	RS1	RS2	RS3	RS4	RS5	RS6	RS7	RS8	Habitat services	HS1	HS2	Provision services	PS1	PS2	PS3	PS4	Carrier services	CS1	CS5	CS6	Information services	IS1	IS2	IS3	IS4	IS5
TH12	Nutrient-rich moist tall herbs		4	4	4	3	5	5	5	4		4	4		2	1	1	3		1	1	1	3	1	1	1	1	2
MD13	Nutrient-poor, extensive moist meadow		3	4	4	3	4	4	4	5		5	4		3	3	3	5		1	1	1	4	2	1	1	1	4
MD14	Nutrient-rich moist and wet meadow		3	4	4	3	4	4	4	3		4	3		3	1	2	2		3	1	1	2	1	1	1	1	1
SB15	Nutrient-poor moist shrub succession		4	3	4	3	5	5	4	4		5	5		2	2	3	3		1	1	1	3	1	2	1	1	1
SB16	Nutrient-rich moist shrub succession		4	3	5	3	5	5	5	3		5	5		3	3	2	2		1	1	1	3	1	1	1	1	2
SB17	Shrubs on semi-moist grassland		4	4	3	3	5	5	5	4		5	5		3	3	2	3		1	1	1	3	3	1	1	1	2
TR18	Chasm woodland		5	4	4	3	4	4	4	3		5	5		3	3	2	3		1	1	1	5	5	3	5	4	4
TR19	Abandoned fruit trees		4	3	3	2	4	4	5	3		5	5		3	3	3	2		1	1	1	5	5	3	5	4	4
TH20	Nutrient-rich tall herbs		3	4	3	2	3	4	4	3		4	3		2	2	1	4		1	1	1	2	1	1	1	1	2
TH21	Nutrient-rich herbs (blooming aspect)		3	3	3	2	3	4	4	3		4	3		2	2	1	2		1	1	1	4	1	1	1	1	2
MD22	Semidry managed meadows		2	3	3	3	4	5	4	4		4	3		2	1	2	3		5	1	1	4	4	2	2	3	3
MD23	Intermediate managed meadows		2	4	3	3	4	5	4	4		4	3		3	1	2	3		5	1	1	3	3	2	2	3	3
MD24	Intensively managed meadows and pastures		2	3	3	2	3	3	3	3		3	2		1	1	1	1		5	1	1	2	1	2	2	2	2

Table 4 includes the second group of classes (classes 12-24) for which the landscape services assessment was carried out. The classes include moist and wet meadows, moist shrubs, tall herbs, chasm woodland, abandoned fruit trees, and meadows subjected to different levels of management.

Regulating services are very well-represented by this group. Unlike the water-related classes, in this group all classes provide medium to very high capacities for most of services, including Soil formation and Soil retention. On the other hand, lower scores were recorded for the service Water supply (from 2 to 3). The highest scores are visible in the Soil formation, Soil retention and Nutrient regulation services, delivered by wet and moist meadows and nutrient-poor moist shrub successions. Lightly and intermediate managed meadows offer higher regulating capacity than intensively managed meadows and pastures. Almost all the classes deliver very high or high capacity for the **Habitat** services. Lower scores were recorded for the intensively managed meadows and pastures. Among the meadows, nutrient-poor extensive moist meadows show the highest capacity for both Refugium and Nursery.

Similarly, to the water-related classes, the group in Table 4 offers low to maximum medium capacity for **Provisioning** services. However, nutrient-poor extensive moist meadows and nutrient-rich tall herbs deliver very high and high capacity for Medicinal resources, respectively.

For the category of the **Carrier** services the open land classes 12-24 have not capacity for the landscape services Transportation and Tourism facilities. The landscape services Cultivation shows capacity within the meadow-related classes exclusively. The semidry meadows as well as intermediate and intensively managed meadows and pastures receive the highest scores (5) whereas the moist meadows show medium capacity (score 3) for the nutrient rich meadows, and no capacity (score 1) for the nutrient poor meadows.

The **Information** services show highest capacities for the chasm woodland and the abandoned fruit trees with scores 3-5 for all **Information** landscape services. Landscape service Aesthetic information shows capacity for all of the open land classes 12-24. Next to the chasm woodland and the abandoned fruit trees, the extensive moist meadows as well as the semidry and intermediate managed meadows are rated especially high (scores 4-5) for their blooming aspects.

Table 5: Capacity matrix of the open land (OL) classes 25-36

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Table 5 includes the third group of classes (classes 25-36) for which the landscape services assessment was carried out. These variegated classes include dry and semi-dry shrubland and fallow land, forest edges, *Pinus nigra* forests on dry grassland, rock fragments and dry-stone walls, field margins, and ruderal vegetation.

Within the **Regulating** services, forest provides the highest capacity for Local Climate Regulation. Soil retention, Soil formation, Nutrient regulation and Pollination are highly represented by this group, especially by nutrient-poor managed meadows and pastures, dry grassland and dry shrubland, and forests like *Pinus nigra* forests. Ruderal Vegetation provides medium capacity for most regulating services, but high capacity for Soil formation.

Almost all the classes deliver very high or high capacity for the **Habitat** services, with semi-dry and dry fallow land of fields and vineyards, species-poor field margins, and ruderal vegetation excluded. In particular, nutrient-poor managed meadows and pastures, semidry and dry grassland, species-rich slopes and margins have very high capacity for Refugium.

Provisioning services are delivered at different potential by this group. While Food shows medium and low scores, Raw materials service is highly provided by forest-related classes. Forest edges also deliver high capacity for Genetic resources and Medicinal resources, together with nutrient-poor managed meadows and pastures and semidry and dry grassland for the latter service.

Landscape service Cultivation has a high capacity in nutrient poor meadows and pastures (score 4) and is the only **Carrier** service addressed by open land classes 25-36.

Except for the ruderal vegetation and species-poor field margins each of the open land classes 25-36 provides capacity for all **Information** services. For the blooming aspects of the nutrient poor meadow, the landscape structuring effect of shrubs, forest edges and stone walls and the particular beauty of the *Pinus nigra* forest on dry grassland high-scores are received in the landscape services Aesthetic information. The accessibility of dry grasslands as well as the shading aspect of the forest edges potentially provide high capacities for landscape services Recreation. The rock fragments and dry-stone wall provide high capacities in landscape services Aesthetic information and Cultural and artistic information. Additionally, they potentially provide rich habitats of interest for landscape services Science and education and have developed over time, therefore providing high capacity for landscape services Spiritual and historic Information.

Table 6: Capacity matrix of open land (OL) classes 37-50

OL Code	Open Land Class	RS1	RS2	RS3	RS4	RS5	RS6	RS7	RS8	Habitat services	HS1	HS2	Provision services	PS1	PS2	PS3	PS4	Carrier services	CS1	CS5	CS6	Information services	IS1	IS2	IS3	IS4	IS5
CU37	Cultivated field	1	1	2	1	1	2	2	2		2	1		5	1	1	1		5	1	1		2	2	2	2	2
CU38	Vineyard	1	1	2	1	2	2	2	2		2	1		4	2	2	1		4	1	1		5	4	5	5	4
CU39	Intensively managed orchard	2	2	3	2	3	4	2	2		2	1		5	2	1	1		5	1	1		2	1	1	1	1
SB40	Nitrophilous and neophytic edges and woodlets	3	2	4	3	3	5	3	3		3	2		2	2	2	2		1	1	1		3	1	1	1	2
SB41	Hedges and shrubs	3	4	4	3	5	5	5	4		4	4		3	3	2	2		1	1	1		4	1	2	2	4
SB42	Shrubs on moist soil	3	4	4	3	5	5	5	4		4	4		3	3	2	3		1	1	1		3	1	2	2	3
TR43	Long-rotation woodland	4	5	5	4	5	5	5	3		4	4		2	4	2	2		1	1	1		4	2	2	2	3
TR44	Hedgerows dominated by trees	4	5	4	4	5	5	5	4		4	4		2	4	2	2		1	1	1		5	3	2	2	4
TR45	Tree rows and single trees	3	3	3	3	3	3	3	4		4	2		2	2	1	1		1	1	1		5	4	5	5	5
TR46	Larch dominated meadows/pastures	3	4	3	3	5	5	4	4		4	3		2	2	2	3		2	1	1		5	5	5	5	5
TR47	Riparian woodland	5	5	5	5	5	5	5	4		5	5		3	4	3	4		1	1	1		5	4	2	3	4
TR48	Extensive orchards	3	4	3	4	5	5	4	5		5	4		4	3	5	3		4	1	1		5	5	5	5	5
TR49	Short-rotation plantations	3	3	3	2	4	4	4	3		3	3		1	4	2	2		1	1	1		1	1	1	1	1
RC50	Cliffs	1	1	1	1	1	2	1	1		4	3		1	1	1	1		1	1	1		5	4	2	3	4

Table 6 includes the fourth group of classes (classes 37-50) for which the landscape services assessment was carried out. Mostly human-managed classes are comprised in this group, such as cultivated fields, vineyards, orchards, hedges, tree rows, plantations, and cliffs. Classes with higher level of naturalness are also included (larch dominated meadows, extensive orchards).

All forest and tree-related classes have intermediate or high capacity for most **Regulating** services. On the other hand, cultivated fields, vineyards, and cliffs deliver no or little regulating capacity. Intensively managed orchards and nitrophilous and neophytic edges and woodlots show high capacity for Soil formation, and the latter for Water regulation as well.

The two Habitat services show high scores for hedges, shrubs, tree rows, and larch dominated meadows, and very high scores for riparian woodland and extensive orchards. Intensively managed classes provide instead low Refugium and Nursery capacity.

Provisioning services are delivered scatterly across the group, with Food showing high scores for cultivated fields, vineyards, and orchards. Woodlands and plantations offer high capacity for Raw materials, whereas extensive orchards offer high capacity for Genetic resources. Medicinal resources are offered at intermediate level by larch dominated meadows, extensive orchards, shrubs on moist soil, and at high level by riparian woodland.

Within the category of Carrier services, only the capacity of landscape services Cultivation is addressed by open land classes 37-50. Cultivated fields, vineyards and orchards provide high capacity (scores 4-5).

For the **Information** Services, vineyards, tree rows and single trees, larch dominated meadows and extensive orchards have high capacity (scores 4-5) for all the services, whereas short-rotation plantations provide no capacity within this service group. Generally, the shrub- and tree-related classes have medium to high capacity for the landscape services Aesthetic information, due their influence on the mosaic display of the landscape in the Wienerwald Biosphere Reserve.

Table 7 includes the fifth group of classes (classes 51-62) for which the landscape services assessment was carried out. Mostly heavily human-managed and rather artificial/transformed classes are comprised in this group, such as cemeteries, waste sites, paved and unpaved open areas, pits and quarries, and gardens or parks.

These classes offer no or little capacity for the **Regulating** services. Medium scores are associated to the less transformed classes only, as gardens or parks, greenery and planting, and cemeteries. These three classes also revealed intermediate or high scores for Pollination.

Habitat services are in general under-represented in this group. Only the above-mentioned three less transformed classes, together with abandoned stone quarries, show intermediate scores for both **Habitat** services.

Similarly, this group delivers no capacity for the **Provisioning** services, if we exclude active gravel pits and quarries for Raw Materials (score 5), and low capacity for Medicinal resources for gardens or parks, and greenery and planting.

The **Carrier** services Transportation and Tourism facilities both show capacity (scores 3-4) in open areas that can be used as parking areas. For the landscape services Tourism facility leisure and sport facilities as well as gardens or parks provide medium capacity (score 3).

The abandoned stone quarries and gravel pits show capacity for all **Information** services (scores 2-3) and especially landscape services Recreation (score 4). Most of the classes within this group provide at least low capacity for the landscape services Recreation; highest are the lawns, cemeteries together with the stone quarries. The medium scores (score 3) in landscape services Recreation for gardens or parks as well as leisure and sport facilities is due to a splitting of the score with landscape services Tourism facilities. Gardens or parks together with greenery and planting are high rated for the landscape services Aesthetic information (score 4-5). Gardens or parks, cemeteries and isolated buildings provide high capacity in landscape services Cultural and artistic and Spiritual and historic Information.

Table 8 includes the sixth group of classes (additional spatial indicators classes 1-18) for which the landscape services assessment was carried out. The table includes all additional spatial indicators classes, i.e. roads, accommodation facilities, gastronomy, museums and castles, religious elements and so on.

The additional spatial indicators are especially important for the groups of **Carrier** and **Information** landscape services, nevertheless the castles and ruins as well as archaeological sites and church towers show low capacity (score 2) for the **Habitat** landscape services, mainly referring to birds of prey and bats inhabiting and breeding in the towers. There are no capacities within the categories of **Regulating** and **Provisioning** services.

Within the **Carrier** Services, the additional spatial indicators classes have no capacity for the landscape services Cultivation. The landscape service Transportation is addressed with high capacity for public transport and primary – tertiary roads (scores 4-5) and medium to low capacity (scores 2-3) for residential roads and trails, which include footpaths and cycling trails. The landscape service Tourism facilities has medium scores for trails as well as look out towers and recreational infrastructure, due to a splitting of scores with the **Information** service Recreation, and high scores (5) for gastronomy and accommodation facilities.

For the **Information** services, the religious buildings, castles and museums show high capacities for all services. The religious elements, including wayside shrines and crosses, and artwork, including monuments and memorials, show similar capacities for similar services. Artwork furthermore provides high capacity (score 5) for landscape service Cultural and artistic information. The landscape service Recreation differs stronger from the other services by valuing trails and viewpoints as well as recreational infrastructure with medium capacity.

The results of the capacity matrix are further integrated into the assessment of capacity intensity within the landscape service capacity maps, described in the next chapter.

5.2 Landscape service capacity maps

The landscape service capacity maps are computed by the integration of the intensity (taken from the capacity matrix values) and the density of the selected landscape elements within the open land of the Wienerwald Biosphere Reserve. According to the INDEC (see chapter 4.8) the maps display clusters ranging from high (deep red) to low (light yellow) capacity of the open land. The colour grey symbolizes both areas of no capacity and the outline of the open land.

Due to limitations in data sources completeness and availability, the three **Carrier** services Habitation, Waste disposal and Energy conversion were dismissed. Furthermore, the recognition of unstandardized data in the open land dataset led to the elimination of data along the “Thermenlinie” of the Wienerwald Biosphere Reserve. Therefore, the maps displayed in the following sections do not represent the whole study region. This bias within the data set of the south-eastern area of the Biosphere Reserve shall be taken into consideration in the examination of the results.

In the following sections the capacity map of each landscape services is represented and described by highlighting (i) the spatial distribution of the service capacity in the study area; (ii) the landscape elements showing the highest capacity in the capacity matrix; and (iii) the three highest ranked clusters according to the INDEC. For the three highest capacity clusters, we provide a table showing the INDEC values, the number of landscape elements within each cluster, the size of each cluster and the distribution of landscape elements per classes within each cluster.

5.2.1 Local climate and air regulation

Figure 11: Local climate and air regulation capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENsity Capacity (INDEC).

Spatial distribution: the service capacity is scatterly distributed. It is almost absent in the NW area between Neulengbach and Tullnerfeld, and very low in the area N of Heiligenkreuz. Large clusters of high capacity are visible in the area of Königstetten and Wolfpassing, at the biosphere reserve border.

Landscape elements: The classes with the highest score (5) according to the capacity matrix are straight, seminatural streams, meandering, seminatural streams, channel/regulated streams, seminatural, nutrient-poor lakes and pools, nutrient-rich pools, artificial standing water connected and disconnected to groundwater, reed beds, chasm woodland, forest edges, and riparian woodland. The classes with score “4” are Periodic/small brooks, Springs, marshes, swamps, nutrient-poor fens, nutrient-rich moist tall herbs, nutrient-poor moist shrub succession, shrubs on semi-moist grassland, abandoned fruit trees, semidry and dry grassland, dry forest edges, *Pinus nigra* forests on dry grassland, long-rotation woodland, hedgerows dominated by trees.

Three clusters with highest INDEC: The first cluster (INDEC 98,61) is located close to Königstetten, in the N of the biosphere reserve, and it includes 66 landscape elements, with a larger percentage of forest edges, hedgerows dominated by trees, and intermediate managed meadows to a fewer extent. The second cluster (INDEC 85,57) is situated E of Liesing, and includes 62 landscape elements, belonging mostly to nutrient-poor extensive moist meadows (20), semidry managed meadows (8) and forest edges (8). Cluster 3 (INDEC 59,69) has a far fewer number of landscape elements (39) and it lies NE of Cluster 1, close to Wolfpassing. It includes landscape elements belonging mostly to forest edges, but also to hedgerows dominated by trees and hedges and shrubs to a minor extent.

Table 9: Three clusters with highest INDEC for Local climate and air regulation. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes’ code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes										
Cluster 1	98,61	66	155.923	TR31	TR44	MD23	TR48	GD56	TH12	MD22	SB17	SB41	TR18	CU39
				17	10	7	6	5	5	4	2	2	2	1
				DR25	MD13	MG36	TR19	TR43						
				1	1	1	1	1						
Cluster 2	85,57	62	112.694	MD13	MD22	TR31	MD14	MD23	GD56	TR44	WT9	MD24	TH12	WT11
				20	8	8	5	5	4	4	4	2	1	1
Cluster 3	59,69	39	98.925	TR31	TR44	SB41	GD56	DR25	MD13	MD23	TH12	TR47	TR48	MD22
				13	5	4	3	2	2	2	2	2	2	1
				SB17										
				1										

5.2.2 Disturbance prevention

Figure 12: Disturbance prevention capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC).

Spatial distribution: The service Disturbance Prevention shows high or very high capacity in most of the open land. However, the large cultivated area in the NW of the reserve, just SW of Asperhofen, presents no service capacity. Intermediate spots of capacity are visible in the area of Tullnerbach, Breitenfurt and Altenglengbach.

Landscape elements: The classes with the highest score (5) according to the capacity matrix are straight, seminatural streams; meandering seminatural streams; periodic/small brooks; long-rotation woodland; hedgerows dominated by trees; riparian woodland. a score “4” was assigned to channel/regulated streams; seminatural, nutrient-poor lakes and pools; nutrient-rich pools; artificial standing water connected to groundwater; springs, marshes, swamps; nutrient-rich moist tall herbs; reed beds; nutrient-rich moist tall herbs; nutrient-poor, extensive moist meadow; nutrient-rich moist and wet meadow; shrubs on semi-moist grassland; chasm woodland, nutrient-rich tall herbs; intermediate managed meadows; dry shrubland on limestone; thermophilous dry shrubland; dry forest edges; forest edges; *Pinus nigra* forests on dry grassland; hedges and shrubs; shrubs on moist soil; larch dominated meadows/pastures; extensive orchards.

Three clusters with highest INDEC: The first cluster (INDEC 136,61) is very large and situated just S of Königstetten and includes 91 landscape elements distributed across 17 classes. The most frequent classes are Forest edges (27), followed by intermediate managed meadows (13) and hedgerows dominated by trees (8). Cluster 2 (INDEC 96,64) lies NE of Breitenfurt bei Wien, in the central area of the Biosphere Reserve and it comprises 67 landscape elements across 15 classes. All meadows are included in this cluster. The most frequent classes are nutrient-poor extensive moist meadows with 20 landscape elements, semidry managed meadows (8), forest edges (8), intermediate managed meadows (7), and hedgerows dominated by trees (7). The third cluster (INDEC 76,28) includes 50 landscape elements and its size is slightly smaller than cluster 2. It is located E of Laaben bei Neulengbach, at the south-western reserve border. The most frequent classes are forest edges, nutrient-poor extensive moist meadows, hedgerows dominated by trees.

Table 10: Three clusters with highest INDEC for Disturbance prevention. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes’ code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes											
Cluster 1	136,61	91	218.083	TR31	MD23	TR44	MD22	SB41	TH12	GD56	TR48	DR25	MD13	MG33	
				27	13	8	6	6	6	5	4	3	2	2	
				SB16	SB17	TR43	MG35	MG36	TR47						
				2	2	2	1	1	1						
Cluster 2	96,64	67	141.744	MD13	MD22	TR31	MD23	TR44	MD14	WT9	MD24	DR25	MG35	SB41	
				20	8	8	7	7	5	4	3	1	1	1	
				TH12	WT11										
				1	1										
Cluster 3	76,28	50	125.645	TR31	MD13	TR44	MD23	MD26	TR45	TR47	WT6	GD57	MD22	TH12	
				13	10	8	4	2	2	2	2	1	1	1	
				TR18	TR43	TR48	TR49								
				1	1	1	1								

5.2.3 Water regulation

Figure 13: Water regulation capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DEnsity Capacity (INDEC)

Spatial distribution: The service capacity for Water regulation presents a similar pattern to Water supply, but with higher capacity altogether. Areas of higher capacity are localized along streams and pools, such as S of Purkersdorf, and W of Liesing, and in the area of Pressbaum (Wienerwald See). Areas of high capacity are also visible in the dense mosaic of landscape elements S of Königstetten. The large north-eastern region of the Biosphere Reserve is dominated by cultivated fields and shows low capacity. Another area with many low, intermediate and high INDEC values is located in N of Alternmarkt an der Triesting.

Landscape elements: The classes with the highest scores ("5" or "4") according to the capacity matrix are straight, seminatural streams, meandering, seminatural streams, periodic/small brooks, seminatural, nutrient-poor lakes and pools, nutrient-rich pools, artificial standing water connected to groundwater, springs, marshes, swamps, nutrient-poor fens, reed beds, nutrient-rich moist tall herbs, nutrient-poor, extensive moist meadow, nutrient-rich moist and wet meadow, nutrient-poor moist shrub succession, nutrient-rich moist shrub succession, chasm woodland, nutrient-poor managed meadows and pastures, forest edges, nitrophilous and neophytic edges and woodlets, hedges and shrubs, shrubs on moist soil, long-rotation woodland, hedgerows dominated by trees, and riparian woodland.

Three clusters with highest INDEC: The first cluster (INDEC 116,8) is located NE of Breitenfurt bei Wien, in the central area of the Biosphere Reserve, almost overlapping cluster 2 of Water supply. It comprises 83 landscape elements, distributed across 15 classes. Nutrient-poor, extensive moist meadows are the most frequent ones, followed by forest edges, semidry managed meadows and cultivated fields. Reed beds are also present in this cluster. Cluster 2 (INDEC 101,45) has 69 landscape elements and a slightly small size than cluster 1. It consists of a great number of classes (19), mostly represented by forest edges, hedgerows dominated by trees, intermediate managed meadows, and extensive orchards. It is situated just S of Königstetten, as cluster 1 of Water supply. The third cluster (INDEC 74,55) is located close to Wöllersdorf, in proximity with the western border. It comprises 49 landscape elements across 16 classes, mostly represented by forest edges, nutrient-poor, extensive moist meadows, and hedgerows dominated by trees. Two nutrient-rich pools occur as well.

Table 11: Three clusters with highest INDEC for Water regulation. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes											
Cluster 1	116,8	83	161.630	MD13	TR31	MD22	CU37	MD23	TR44	MD14	MD24	GD56	WT9	DR25	
				21	10	8	7	7	7	5	5	4	4	1	
				MG35	SB41	TH12	WT11								
				1	1	1	1								
Cluster 2	101,45	69	155.165	TR31	TR44	MD23	TR48	GD56	TH12	MD22	CU38	DR25	SB41	TR18	
				16	10	7	6	5	5	4	2	2	2	2	
				CU39	MD13	MG36	SB17	SB40	TR19	TR43	TR47				
				1	1	1	1	1	1	1	1				
Cluster 3	74,55	49	122.185	TR31	MD13	TR44	MD23	MD24	MD26	TR18	TR43	TR47	WT6	GD56	
				13	8	7	3	2	2	2	2	2	2	1	
				GD57	TH12	TR46	TR48	TR49							
				1	1	1	1	1							

5.2.4 Water supply

Figure 14: Water supply capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC).

Spatial distribution: the service capacity is expressed through rather small clusters, with areas of highest capacity localized along streams and pools, such as S of Purkersdorf, and W of Liesing, and in the area of Pressbaum (Wienerwald See). Areas of high capacity are also visible in the dense mosaic of landscape elements S of Königstetten. The large north-eastern region of the Biosphere Reserve is dominated by cultivated fields and shows no capacity. Another area with many low, intermediate and high INDEC values is located in N of Alternmarkt an der Triesting.

Landscape elements: The classes with the highest scores according to the capacity matrix are straight, seminatural streams, meandering, seminatural streams, channel/regulated streams, seminatural, nutrient-poor lakes and pools, nutrient-rich pools, artificial standing water connected to groundwater, springs, marshes, swamps, nutrient-poor fens, riparian woodland. A “4” was assigned to periodic/small brooks, reed beds, forest edges, long-rotation woodland, hedgerows dominated by trees, and extensive orchards.

Three clusters with highest INDEC: The first cluster (INDEC 86,6) is just SE of Königstetten and includes 60 landscape elements distributed across 15 classes. The most frequent classes are forest edges, followed by intermediate managed meadows and hedgerows dominated by trees. Cluster 2 (INDEC 83,09) is a mosaic of meadows and lies W of Liesing; it has 60 landscape elements distributed across 11 classes. The most frequent classes are nutrient-poor extensive moist meadows, semidry managed meadows, and forest edges. Cluster 3 (INDEC 56,24) is situated in the southern area of the Biosphere Reserve, close to Obertriesting, in a mosaic of managed meadows, tree rows and hedges. It includes 40 landscape elements distributed across 10 classes, whose most frequent are forest edges, tree rows and single trees, and intermediate and intensively managed meadows.

Table 12: Three clusters with highest INDEC for Water supply. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes										
Cluster 1	86,6	60	127.218	TR31	MD23	TR44	MD22	TH12	SB41	MD13	SB17	TR43	TR48	DR25
				18	11	7	4	4	3	2	2	2	2	1
				MG33	MG35	MG36	SB16							
				1	1	1	1							
Cluster 2	83,09	60	110.431	MD13	MD22	TR31	MD23	TR44	MD14	WT9	MD24	SB41	TH12	WT11
				20	8	7	6	6	5	4	1	1	1	1
Cluster 3	56,24	40	77.658	TR31	TR45	MD23	MD24	MD14	TH12	TR43	TR44	MG35	TH20	
				10	6	5	5	4	3	3	2	1	1	

5.2.5 Soil retention

Figure 15: Soil retention capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC).

Spatial distribution: the service Soil retention shows a moderate to high capacity in most of the open areas of the reserve. The only excluded area is the large cultivated sector in the NW of the Biosphere Reserve. Very large clusters are visible along the northern border and in the municipality of Vienna as well. Streams and pools are not providers of the service, and for this reason the areas rich in water-related classes show lower capacity.

Landscape elements: The classes with the highest scores ("4" or "5") according to the capacity matrix are reed beds, nutrient-rich moist tall herbs, nutrient-poor, extensive moist meadow, nutrient-rich moist and wet meadow, nutrient-poor moist shrub succession, nutrient-rich moist shrub succession, shrubs on semi-moist grassland, chasm woodland, abandoned fruit trees, semidry managed meadows, intermediate managed meadows, nutrient-poor managed meadows and pastures, dry shrubland on limestone, thermophilous dry shrubland, semidry and dry grassland, dry forest edges, forest edges, *Pinus nigra* forests on dry grassland, hedges and shrubs, shrubs on moist soil, long-rotation woodland, hedgerows dominated by trees, larch dominated meadows/pastures, riparian woodland, extensive orchards, and short-rotation plantations.

Three clusters with highest INDEC: The first cluster (INDEC 199,43) is a dense mosaic of LE located S of Königstetten. It includes a very high number of LE (129), distributed across 22 classes. The most frequent five are forest edges, intermediate managed meadows, reed beds, semidry managed meadows, gardens or parks, and hedges and shrubs. Cluster 2 (INDEC 102,84) lies just E of cluster 1, but it is much smaller and it has a far fewer number of LE (66). The most frequent classes are extensive orchards, vineyards, semi-dry and dry fallow land of fields and vineyards, species-rich slopes and margins, and intermediate managed meadows. The third cluster (INDEC 97,38) is located in the southern area, N of Altenmarkt an der Triesting, close to the little settlement Pankrazi. This cluster is way less variegated than the other two, with 62 landscape elements distributed across 11 classes. The three most frequent are hedgerows dominated by trees, tree rows and single trees, and intermediate managed meadows.

Table 13: Three clusters with highest INDEC for Soil retention. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes										
Cluster 1	199,43	129	336.765	TR31	MD23	TR44	MD22	GD56	SB41	TH12	TR48	SB17	DR25	MD13
				32	17	11	10	8	7	7	7	5	3	3
				SB16	TR45	CU38	MG33	TR18	TR43	CU39	MG35	MG36	TR19	TR47
				3	3	2	2	2	2	1	1	1	1	1
Cluster 2	102,84	66	176.185	TR48	CU38	DR25	MG33	MD23	TR44	GD56	MD22	SB41	TR18	TR31
				8	6	6	6	5	5	3	3	3	3	3
				MD24	MG35	SB40	TR43	TR49	DR30	MD26	SB16	TH20	TR19	
				2	2	2	2	2	1	1	1	1	1	
Cluster 3	97,38	62	169.201	TR44	TR45	MD23	TR31	MD24	TR48	MD13	GD56	MG35	MD22	WT9
				12	12	8	8	6	6	4	2	2	1	1

5.2.6 Soil formation

Figure 16: Soil formation capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DEnsity Capacity (INDEC).

Spatial distribution: the service is more concentrated in the northern border of the Reserve, between the areas S of Baumgarten am Tullnerfeld and S of Königstetten. Very large clusters can be detected in this area, whereas the remainder sector of the open land show moderate to high INDEC and small clusters. Areas of lower capacity are visible along the central sector of the Biosphere Reserve. Another important concentration of clusters is located in the southern sector, NE of Altenmarkt an der Triesting.

Landscape elements: Most of the open land classes have high capacity for this service. The classes with the highest scores ("4" or "5") according to the capacity matrix are springs, marshes, swamps, nutrient-poor fens, reed beds, nutrient-rich moist tall herbs, nutrient-poor, extensive moist meadow, nutrient-rich moist and wet meadows, nutrient-poor moist shrub succession, nutrient-rich moist shrub succession, shrubs on semi-moist grassland, chasm woodland, abandoned fruit trees, nutrient-rich tall herbs, nutrient-rich herbs (blooming aspect), semidry managed meadows, intermediate managed meadows, semi-dry and dry fallow land of fields and vineyards, nutrient-poor managed meadows and pastures, dry shrubland on limestone, thermophilous dry shrubland, semidry and dry grassland, dry forest edges, forest edges, *Pinus nigra* forests on dry grassland, species-rich slopes and margins, ruderal vegetation, intensively managed orchard, nitrophilous and neophytic edges and woodlets, hedges and shrubs, shrubs on moist soil, long-rotation woodland, hedgerows dominated by trees, larch dominated meadows/pastures, riparian woodland, extensive orchards, and short-rotation plantations.

Three clusters with highest INDEC: The three clusters, all very large and very rich in landscape elements, are located one next to the other, between S of Königstetten and S of Tulbing. The first cluster (INDEC 275,43) is an extremely large (m² 354.956) and dense mosaic of landscape elements, situated S of Königstetten. It includes a very high number of landscape elements (130), distributed across 23 classes. The most frequent five are forest edges, intermediate managed meadows, hedgerows dominated by trees, semidry managed meadows, and gardens or parks. Cluster 2 (INDEC 204,23) lies S of Tulbing and it includes 120 landscape elements across 23 classes, like cluster 1, but has a small size. The most frequent five are vineyards, intermediate managed meadows, forest edges, hedgerows dominated by trees, and intensively managed meadows and pastures. these classes are evenly distributed. The third cluster (INDEC 186,95) is also a very variegated, large landscape mosaic, with 102 landscape elements included into 22 classes. It is located between cluster 1 and cluster 2. The most frequent five are semi-dry and dry fallow land of fields and vineyards, vineyards, hedges and shrubs, extensive orchards, and intermediate managed meadows.

Table 14: Three clusters with highest INDEC for Soil formation. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes											
Cluster 1	275,43	130	354.956	TR31	MD23	TR44	MD22	GD56	SB41	TH12	TR48	SB17	DR25	MD13	
				32	17	11	10	8	7	7	7	5	3	3	
				SB16	TR45	CU38	MG33	TR18	TR43	CU39	MG35	MG36	SB40	TR19	TR47
				3	3	2	2	2	2	1	1	1	1	1	1
Cluster 2	204,23	120	320.152	CU38	MD23	TR31	TR44	MD24	MD22	CU37	DR25	SB41	TR48	GD56	
				14	13	11	10	9	8	7	7	7	7	5	
				MG33	MD13	MD26	SB40	MG35	AR55	TH12	TR18	TR19	TR43	TR45	TR49
				4	3	3	3	2	1	1	1	1	1	1	1
Cluster 3	186,95	102	309.003	DR25	CU38	SB41	TR48	MD23	MG33	TR31	TR44	GD56	MD22	MG35	
				15	10	9	9	8	6	6	5	4	4	4	
				TR18	TR43	MD24	DR30	SB40	TR49	CU37	MD26	SB16	TH20	TR19	
				4	4	3	2	2	2	1	1	1	1	1	

5.2.7 Nutrient regulation

Figure 17: Nutrient regulation capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC)

Spatial distribution: the service Nutrient regulation is very well represented in the Biosphere Reserve, and more concentrated in the northern border of the Reserve, as for most of the other regulating services. Large clusters are visible between the areas S of Baumgarten am Tullnerfeld and S of Wolfpassing, close to the northern Biosphere Reserve border. Other high INDEC values were detected in the most northern corner of the Biosphere Reserve, in the proximity of Höflein an der Donau. Other high capacity areas can be detected between Klosterneuburg and the municipality of Vienna. More important concentrations of clusters are located in the southern sector, NE of Altenmarkt an der Triesting, and in the proximity of Laaben, in the very western sector of the reserve.

Landscape elements: More than half of the open land classes (36) have a service capacity of either “4” or “5” according to the capacity matrix. They are straight, seminatural streams meandering seminatural streams, channel/regulated streams, periodic/small brooks, seminatural, nutrient-poor lakes and pools, artificial standing water connected to groundwater, springs, marshes, swamps, nutrient-poor fens, reed beds, nutrient-rich moist tall herbs, nutrient-poor, extensive moist meadow, nutrient-rich moist and wet meadow, nutrient-poor moist shrub succession, nutrient-rich moist shrub succession, shrubs on semi-moist grassland, chasm woodland, abandoned fruit trees, nutrient-rich tall herbs, nutrient-rich herbs (blooming aspect), semidry managed meadows, intermediate managed meadows, nutrient-poor managed meadows and pastures, dry shrubland on limestone, thermophilous dry shrubland, semidry and dry grassland, dry forest edges, forest edges, *Pinus nigra* forests on dry grassland, hedges and shrubs, shrubs on moist soil, long-rotation woodland, hedgerows dominated by trees, larch dominated meadows/pastures, riparian woodland, extensive orchards, and short-rotation plantations.

Three clusters with highest INDEC: The three clusters are located all on the northern border of the Biosphere Reserve, between S of Freundorf and S of Königstetten. They are large and very rich in landscape elements, especially the first cluster, which has a double size compared to the other two. The first cluster (INDEC 198,78) is located S of Königstetten, it includes 129 landscape elements across 22 classes. Most frequent classes are forest edges (32), followed by 17 landscape elements belonging to intermediate managed meadows, hedgerows dominated by trees (11) and semidry managed meadows (10). Cluster 2 (INDEC 125,2) lies S of Freundorf, comprises 90 landscape elements distributed across 17 classes. The three most frequent classes are species-rich slopes and margins, species-poor field margins, and semi-dry and dry fallow land of fields and vineyards. These classes are evenly distributed. The third cluster (INDEC 115,9) is located in contiguity with cluster 2, thus guaranteeing high service connectivity. It is a very variegated mosaic, consisting of 79 landscape elements distributed across 17 classes, like cluster 2. The most frequent class is cultivated fields (13), followed by intermediate managed meadows (9), species-poor field margins (9) and semi-dry and dry fallow land of fields and vineyards (6). The landscape elements are evenly distributed across the classes.

Table 15: Three clusters with highest INDEC for Nutrient regulation. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC	N LE	Area (m ²)	Distribution of LE classes												
Cluster 1	198,78	129	333.698	TR31	MD23	TR44	MD22	GD56	SB41	TH12	TR48	SB17	DR25	MD13		
				32	17	11	10	8	7	7	7	5	3	3		
				SB16	TR45	CU38	MG33	TR18	TR43	CU39	MG35	MG36	TR19	TR47		
				3	3	2	2	2	2	1	1	1	1	1		
Cluster 2	125,2	90	168.310	MG33	MG35	DR25	CU37	MD23	TR44	CU38	TR48	MD24	SB41	TR18		
				14	14	12	10	7	7	6	6	3	3	2		
				GD56	MD22	SB17	TR43	TR45	TR49							
				1	1	1	1	1	1							
Cluster 3	115,9	79	176.428	CU37	MD23	MG35	DR25	TR44	TR48	CU38	SB41	TR49	GD56	MD22		
				13	9	9	6	6	6	5	5	4	3	3		
				TR45	MD24	MG33	SB42	TH12	TR19							
				3	2	2	1	1	1							

5.2.8 Pollination

Figure 18: Pollination capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC)

Spatial distribution: the service capacity is generally very high in the reserve, and more concentrated in the N of the Biosphere Reserve, particularly on the reserve border in the proximity of Baumgarten am Tullnerfeld, Königstetten, Klosterneuburg, and Döbling in the municipality of Vienna. Other significant areas for Pollination can be found in the southern part of the Biosphere Reserve, in the very large mosaic of meadows and fields just N of Altenmarkt an der Triesting.

Landscape elements: The classes with the highest score according to the capacity matrix are Nutrient-poor extensive moist meadow, Nutrient-poor managed meadows and pastures, Species-rich slopes and margins, and Extensive orchards.

The classes with the score “4” according to the capacity matrix are nutrient-poor fens, nutrient-rich moist tall herbs, nutrient-poor moist shrub succession, shrubs on semi-moist grassland, semidry managed meadows, intermediate managed meadows, dry shrubland on limestone, thermophilous dry shrubland, semidry and dry grassland, dry forest edges, forest edges, *Pinus nigra* forests on dry grassland, hedges and shrubs, shrubs on moist soil, hedgerows dominated by trees, tree rows and single trees, larch dominated meadows/pastures, riparian woodland, gardens or parks, and greenery and planting.

Three clusters with highest INDEC: The first cluster (INDEC 183,68) is a very large group of 121 landscape elements, distributed across 24 classes, and it is located close to Königstetten, on the reserve northern border. the three most frequent classes are forest edges (30 landscape elements), followed by intermediate managed meadows (17) and hedgerows dominated by trees (11). other frequent classes are gardens or parks, and semidry managed meadows. Cluster 2 (INDEC 132,7) lies quite close to Cluster 1, in the area of Baumgarten am Tullnerfeld. It includes 91 landscape elements, distributed in 17 classes. The most frequent class is cultivated fields, due to their density and small sizes in this cluster. Second most frequent class is Species-poor field margin, followed by intermediate managed meadow. The third cluster (INDEC 108,81) is again a large cluster including 73 landscape elements. Of these, 25 belong to tree rows and single trees, 14 to hedgerows dominated by trees, and 7 to semidry managed meadows. Other classes present in the cluster are extensive orchards, hedgerows dominated by trees, and forest edges. This cluster is located in the southern part of the Biosphere Reserve, in the proximity of Groisbach.

Table 16: Three clusters with highest INDEC for Pollination. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes’ code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes											
Cluster 1	183,68	121	299.734	TR31	MD23	TR44	GD56	MD22	TH12	TR48	SB41	DR25	SB16	CU38	MD13
				30	17	11	9	9	7	7	6	3	3	2	2
				MG33	SB17	TR18	TR43	CU39	MG35	MG36	SB40	TR19	TR45	TR47	
				2	2	2	2	1	1	1	1	1	1	1	
Cluster 2	132,7	91	199.398	CU37	MG35	MD23	TR48	TR44	CU38	DR25	GD56	SB41	MG33	TR49	
				15	10	9	8	7	6	6	5	5	4	4	
				MD22	MD24	TR45	SB42	TH12	TR43						
				3	3	3	1	1	1						
Cluster 3	108,81	73	171.236	TR45	TR44	MD22	MD24	TR31	TR48	CU37	DR29	MD23	MG33	DR25	
				25	14	7	5	4	4	3	2	2	2	1	
				MD13	MD26	MG35	SB41								
				1	1	1	1								

5.2.9 Refugium

Figure 19: Refugium capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC)

Spatial distribution: Similarly to Pollination, the service capacity is generally very high and very spread in the reserve, and more concentrated in the N of the Biosphere Reserve, particularly on the reserve border in the proximity of Baumgarten am Tullnerfeld, Königstetten, Klosterneuburg, and Döbling in the municipality of Vienna. Other significant areas for Pollination can be found in the southern part of the Biosphere Reserve, in the very large mosaic of meadows and fields just N of Altenmarkt an der Triesting.

Landscape elements: The classes with the highest score according to the capacity matrix are springs, marshes, swamps; nutrient-poor fens; reed beds; nutrient-poor, extensive moist meadow; nutrient-poor moist shrub succession; nutrient-rich moist shrub succession; shrubs on semi-moist grassland; chasm woodland; abandoned fruit trees; nutrient-poor managed meadows and pastures; forest edges; riparian woodland; extensive orchards. The classes with the score “4” according to the capacity matrix are straight, seminatural streams, meandering seminatural streams, periodic/small brooks, seminatural nutrient-poor lakes and pools, nutrient-rich moist tall herbs; nutrient-rich moist and wet meadow; nutrient-rich tall herbs, nutrient-rich herbs (blooming aspect); semidry managed meadows; intermediate managed meadows; dry shrubland on limestone; thermophilous dry shrubland; semidry and dry grassland; dry forest edges; *Pinus nigra* forests on dry grassland; species-rich slopes and margins; rock fragments and dry-stone walls; hedges and shrubs; shrubs on moist soil; long-rotation woodland; hedgerows dominated by trees; tree rows and single trees; larch dominated meadows/pastures; cliffs.

Three clusters with highest INDEC: The first cluster (INDEC 205,1) is very similar to the cluster 1 of Pollination: it is located close to Königstetten, and it includes 131 landscape elements across a very large area. The most frequent classes out of 23 are forest edges (32 landscape elements), followed by Intermediate managed meadows (17) and Hedgerows dominated by trees (11). Other frequent classes are semidry managed meadows and gardens or parks. Cluster 2 (INDEC 137,16) lies E of Freundorf in the norther part of the reserve, and it includes 94 landscape elements across 18 classes, whose most frequent are cultivated fields, Intermediate managed meadows, species-poor field margins, Hedgerows dominated by trees and extensive orchards. The third cluster (INDEC 135,71) is located in the most northern sector of the Biosphere Reserve, just S of Höflein an der Donau. It comprises 88 landscape elements across 25 classes. It is more diverse than the other two clusters, as it includes more classes, which are also more evenly distributed. The most frequent are forest edges, extensive orchards and semidry managed meadows. Other classes are gardens or parks, Semi-dry and dry fallow land of fields and vineyards, chasm woodland, and Hedgerows dominated by trees. An additional spatial indicators class (religious buildings) is present as well.

Table 17: Three clusters with highest INDEC for Refugium. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes											
Cluster 1	205,1	131	354.338	TR31	MD23	TR44	MD22	GD56	SB41	TH12	TR48	SB17	DR25	MD13	SB16
				32	17	11	10	9	7	7	7	5	3	3	3
				TR45	CU38	MG33	TR18	TR43	CU39	MG35	MG36	SB40	TR19	TR47	
				3	2	2	2	2	1	1	1	1	1	1	
Cluster 2	137,16	94	206.362	CU37	MD23	MG35	TR44	TR48	CU38	DR25	GD56	SB41	MG33	TR49	
				15	10	10	8	8	6		5	5	4	4	
				MD22	MD24	TR45	SB42	TH12	TR19	TR43					
				3	3	3	1	1	1	1					
Cluster 3	135,71	88	228.138	TR31	TR48	MD22	GD56	DR25	TR18	TR44	DR29	MD26	MD24	REL9	SB16
				16	15	12	7	4	4	4	3	3	2	2	2
				TR19	AR59	CU38	CU39	GD57	GD58	MD23	SB17	SB41	TH12	TR43	TR45
				2	1	1	1	1	1	1	1	1	1	1	1
				TR47											

5.2.10 Nursery

Figure 20: Nursery capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC)

Spatial distribution: The second **Habitat** service, Nursery, has a similar capacity distribution to Refugium, although much more attenuated. It is more concentrated in the N of the Biosphere Reserve, particularly on the reserve border in the proximity of Baumgarten am Tullnerfeld, Königstetten, Klosterneuburg, and Döbling in the municipality of Vienna. Other areas of high capacity can be visualized in the area of Glashütte.

Landscape elements: The classes with the highest scores according to the capacity matrix are straight seminatural streams, meandering seminatural streams, seminatural nutrient-poor lakes and pools, nutrient-rich pools, reed beds, nutrient-poor moist shrub succession, nutrient-rich moist shrub succession, shrubs on semi-moist grassland, chasm woodland, abandoned fruit trees, riparian woodland. A “4” was assigned to periodic/small brooks, artificial standing water connected to groundwater, artificial pools disconnected from groundwater, springs, marshes, swamps, nutrient-poor fens, nutrient-rich tall herbs, nutrient-poor extensive moist meadow; dry shrubland on limestone; thermophilous dry shrubland; semidry and dry grassland; forest edges, dry forest edges; *Pinus nigra* forests on dry grassland; species-rich slopes and margins; rock fragments and dry-stone walls; hedges and shrubs; shrubs on moist soil; long-rotation woodland; hedgerows dominated by trees; extensive orchards.

Three clusters with highest INDEC: The first cluster (INDEC 140,32) is also for this service located close to Königstetten, but it is smaller and with fewer landscape elements compared to Refugium and Pollination. It includes 95 landscape elements, whose most frequent classes out of 18 are forest edges (27 landscape elements), followed by Intermediate managed meadows (15) and hedgerows dominated by trees (11). Other frequent classes are semidry managed meadows (6), nutrient-rich moist tall herbs (6) and extensive orchards (6). Similarly to Refugium, Cluster 2 (INDEC 100,63) lies E of Freundorf in the norther part of the reserve, and it includes 68 landscape elements across 20 classes, whose most frequent are extensive orchards (14), semidry managed meadows (10), forest edges (10) and gardens or parks. The third cluster (INDEC 96,39) is situated NE of Breitenfurt, in the central area of the Biosphere Reserve, and it comprises 68 landscape elements like the second cluster. It is, however, smaller. The most frequent classes are nutrient-poor extensive moist meadows, intermediate managed meadows, and semidry managed meadows. A wetland is also included in the cluster (reed beds).

Table 18: Three clusters with highest INDEC for Nursery. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes										
Cluster 1	140,32	95	216.714	TR31	MD23	TR44	MD22	TH12	TR48	GD56	SB17	SB41	DR25	MD13
				27	15	11	6	6	6	5	3	3	2	2
				TR18	TR43	MG33	MG35	MG36	SB16	TR19				
				2	2	1	1	1	1	1				
Cluster 2	100,63	68	156.051	TR48	MD22	TR31	GD56	DR25	TR18	DR29	MD26	REL9	TR19	TR44
				14	10	10	5	4	4	3	3	2	2	2
				GD58	MD23	SB16	SB17	SB41	TH12	TR43	TR45	TR47		
				1	1	1	1	1	1	1	1	1		
Cluster 3	96,39	68	135.761	MD13	TR31	MD22	MD23	TR44	MD14	GD56	WT9	MD24	SB41	TH12
				21	9	8	6	6	5	4	4	2	1	1
				WT11										

5.2.11 Food

Figure 21: Food capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC)

Spatial distribution: the service capacity is more concentrated and also very high in the NW area of the Biosphere Reserve, where a very large cultivated area occurs. The cultivated fields in this region have an extremely large size compared to the other landscape elements in the Biosphere Reserve. Other single clusters can be seen in the central area of the reserve, N of Heiligenkreuz, where some very large cultivated fields occur.

Landscape elements: The classes with the highest scores according to the capacity matrix are cultivated fields (5), intensively managed orchards (5), nutrient-rich pools (4), vineyards (4), and extensive orchards (4).

Three clusters with highest INDEC: The first cluster (INDEC 80,84) is located S of Baumgarten am Tullnerfeld, it comprises 59 landscape elements across 13 classes. The most frequent classes are extensive orchards, cultivated fields, and to a minor extent semi-dry and dry fallow land of fields and vineyards, and intermediate managed meadows. Cluster 2 (INDEC 78,17) lies very close to the first cluster, has 51 landscape elements distributed mostly across species-rich slopes and margins, cultivated fields, semi-dry and dry fallow land of fields and vineyards, and intermediate managed meadows. The third cluster (INDEC 68,75) is located NE of Breitenfurt, in the central area of the Biosphere Reserve, like the third cluster of Nursery. It comprises 48 landscape elements across 12 classes. Most of them belong to nutrient-poor extensive moist meadows (15), cultivated fields (7), semidry managed meadows (7) and forest edges (7).

Table 19: Three clusters with highest INDEC for Food. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes											
Cluster 1	80,84	59	104.457	TR48	CU37	DR25	MD23	TR18	CU38	SB41	TR44	DR29	MG33	MD22	
				12	11	4	4	4	3	3	3	2	2	1	
				TR19	TR43										
				1	1										
Cluster 2	78,17	51	129.927	MG33	CU37	DR25	CU38	TR48	TR44	MD23	SB17	SB41	TR18		
				10	9	7	6	6	4	3	1	1	1		
Cluster 3	68,75	48	99.225	MD13	CU37	MD22	TR31	TR44	MD14	MD23	WT9	DR25	SB41	TH12	WT11
				15	7	7	7	6	5	5	3	1	1	1	1

5.2.12 Raw materials

Figure 22: Raw materials capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DEnsity Capacity (INDEC)

Spatial distribution: the service capacity for Raw Materials is generally very low represented in the Biosphere Reserve (very small clusters and low INDEC values), in comparison with the other services. Some areas of higher capacity can be seen N and NE of the reserve, close to the border. Other single clusters or isolated clusters are visible in the quarry E of Gaaden and close to Altenmarkt an der Triesting, respectively.

Landscape elements: Only one class received a “5” according to the capacity matrix: active gravel pits and quarries. A “4” was assigned to dry forest edges, forest edges, *Pinus nigra* forests on dry grassland, long-rotation woodland, hedgerows dominated by trees, riparian woodland, and short-rotation plantations.

Three clusters with highest INDEC: The first cluster (INDEC 32,25) is located in the SE area of the Biosphere Reserve, in the area of Baden, and it comprises 25 landscape elements. Most of them (14) belong to semidry and dry grassland, then to *Pinus nigra* forests on dry grassland, to dry forest edges and to forest edges. Cluster 2 (INDEC 24,02) is more variegated and larger than Cluster 1, including 6 classes but 16 landscape elements only, whose most frequent belong to Forest edges and Nutrient-poor extensive moist meadows. It is located in the proximity of Wöllersdorf, N of Glashütte. The third cluster (INDEC 22,98) is located N of Altenmarkt an der Triesting and it comprises 4 classes only, with one clearly dominating the others. These are: hedgerows dominated by trees (most frequent with 10 landscape elements), forest edges, tree rows and single trees, and nutrient-poor extensive moist meadows.

Table 20: Three clusters with highest INDEC for Raw materials. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes					
Cluster 1	32,25	25	34.647	DR29	TR32	DR30	TR31		
				14	5	3	3		
Cluster 2	24,02	16	38.347	TR31	MD13	TR44	TR47	TR18	TR49
				6	4	2	2	1	1
Cluster 3	22,98	15	38.158	TR44	TR31	TR45	MD13		
				10	2	2	1		

5.2.13 Genetic resources

Figure 23: Genetic resources capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DEnsity Capacity (INDEC)

Spatial distribution: The service capacity for Genetic Resources is generally very low represented in the Biosphere Reserve (very small clusters and low INDEC values), in comparison with the other services, similarly to Raw Materials. Areas of high capacity can be seen in the N of the Biosphere Reserve, close to Höflein an der Donau, and in the eastern area of the Biosphere Reserve, W of Hietzing. Another area characterized by intermediate and high capacity is around Laaben, in the western part of the Biosphere Reserve.

Landscape elements: The class with the highest score according to the capacity matrix is Extensive orchards, while Forest edges were assigned a “4”.

Three clusters with highest INDEC: The first cluster (INDEC 30,18) is almost the same highlighted in Raw Materials. It is located in the SE area of the Biosphere Reserve, in the area of Baden, and it comprises 25 landscape elements. Most of them (14) belong to semidry and dry grassland, then to *Pinus nigra* forests on dry grassland, to dry forest edges and to semidry managed meadows. Cluster 2 (INDEC 29,59) lies NE of Breitenfurt, in the central area of the Biosphere Reserve. It incorporates 23 landscape elements across 6 classes, whose most frequent are Nutrient-poor extensive moist meadows. Other classes represented are nutrient-rich moist and wet meadows, semidry managed meadows, and intermediate managed meadows. The third cluster (INDEC 26,27) is located on the slopes of Döbling, in the municipality of Vienna. It includes 21 landscape elements nicely distributed across 7 classes. 9 are part of cultivated fields, 3 of nitrophilous and neophytic edges and woodlets, 3 of forest edges, and 2 of hedges and shrubs. Other classes present are nutrient-rich moist shrub succession, extensive orchards and hedgerows dominated by trees.

Table 21: Three clusters with highest INDEC for Genetic resources. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes’ code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes						
Cluster 1	30,18	25	24.781	DR29	TR32	DR30	MD22	TR31		
				14	4	3	2	2		
Cluster 2	29,59	23	31.499	MD13	MD14	MD22	MD23	TR31	TR44	
				11	5	3	2	1	1	
Cluster 3	26,27	21	25.216	CU38	SB40	TR31	SB41	TR48	SB16	TR44
				9	3	3	2	2	1	1

5.2.14 Medicinal resources

Figure 24: Medicinal resources capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DEnsity Capacity (INDEC)

Spatial distribution: The service capacity is nicely spread across the Biosphere Reserve, with the exclusion of the large cultivated fields in the NW. Areas of higher capacity are visible between Hietzing and Liesing, and in the very northern sector, close to the reserve border. A large single cluster can be identified close to Sittendorf, in the central sector of the reserve.

Landscape elements: The classes with the highest scores according to the capacity matrix are nutrient-poor fens (4), nutrient-poor extensive moist meadows (5), nutrient-poor managed meadows and pastures (5), nutrient-rich tall herbs (4), semidry and dry grassland (5), forest edges (4), species-rich slopes and margins (4), riparian woodland (4).

Three clusters with highest INDEC: The first cluster (INDEC 61,81) is located NE of Breitenfurt, in the central area of the Biosphere Reserve. It includes 45 landscape elements, mostly Nutrient-poor extensive moist meadows (16), but also semidry managed meadows (8), Nutrient-rich moist and wet meadows (5), and intermediate managed meadows (5). Cluster 2 (INDEC 46,99) lies between St. Andrä and Wolfpassing, on the reserve northern border. It has 34 landscape elements, well distributed across 13 classes. Forest edges are the most frequent, then gardens or parks, nutrient-rich moist tall herbs and chasm woodland. The third cluster (INDEC 46,88) is located S of Wöllersdorf, at the ESE border. It includes 31 landscape elements across 10 classes, whose three most frequent are forest edges, then nutrient-poor extensive moist meadows, intermediate managed meadows, and hedgerows dominated by trees.

Table 22: Three clusters with highest INDEC for Medicinal resources. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes									
Cluster 1	61,81	45	80.364	MD13	MD22	MD14	MD23	TR31	TR44	WT9	TH12	WT11	
				16	8	5	5	3	3	3	1	1	
Cluster 2	46,99	34	62.096	TR31	GD56	TH12	TR18	TR44	MD22	MD23	SB41	DR25	DR29
				8	5	4	3	3	2	2	2	1	1
				MD13	SB16	TR48							
				1	1	1							
Cluster 3	46,88	31	75.958	TR31	MD13	MD23	TR44	MD26	TR47	GD57	MD22	TR18	TR49
				10	7	3	3	2	2	1	1	1	1

5.2.15 Cultivation

Figure 25: Cultivation capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC)

Spatial distribution: the service is more concentrated in the N and NW area, with a higher aggregation of dense clusters in the N area in the surrounding of Tullnerfeld and bordering the biosphere. Isolated spots ("single clusters") of high capacity were also revealed in the area N of Heiligenkreuz, in the southern region of the biosphere, and in the area E of Asperhofen and NE of Neulengbach.

Landscape elements: The classes with the highest scores according to the capacity matrix are cultivated fields (5), semidry managed meadows (5), intermediate managed meadows (5), intensively managed meadows and pastures (5), intensively managed orchards (5), nutrient-poor managed meadows and pastures (4), vineyards (4), extensive orchards (4).

Three clusters with highest INDEC: The first cluster (INDEC 119,87) is located close to Freundorf and it includes a mosaic of very dense small-sized and medium-sized polygons mainly represented by cultivated fields, extensive orchards, vineyards and semidry to intensively managed meadows and pastures. Cluster 2 (INDEC 60,39) lies west of cluster 1 in Baumgarten am Tullnerfeld and incorporates extensive orchards, cultivated fields, vineyards and intensive meadows and pastures. The third cluster (INDEC 41,41) is east of cluster 1 in Klein-Staasdorf and includes cultivated fields, intermediate and intensive managed meadows, vineyards and extensive orchards.

Table 23: Three clusters with highest INDEC for Cultivation. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes						
Cluster 1	119,87	67	252.804	CU37	TR48	CU38	MD22	MD23	MD24	MD26
				25	15	12	6	5	3	1
Cluster 2	60,39	38	107.049	TR48	CU37	CU38	MD24			
				11	9	9	9			
Cluster 3	41,41	24	83.247	CU37	MD23	CU38	TR48	MD24		
				9	5	4	4	2		

5.2.16 Transportation

Figure 26: Transportation capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DEnsity Capacity (INDEC)

Spatial distribution: In the landscape services Transportation high capacity clusters show the location of main roads within the Biosphere Reserve. In the south, Alland is a crossing point of the A21 (Wiener Außenring Autobahn) with the B11 Mödlinger Straße and B210 Badner Straße. In the centre of the Biosphere Reserve, going through Altlenzbach and Pressbaum, the A1 Westautobahn and the B13 Laaber Straße are visible running towards Vienna above and below the Lainzer Tiergarten. The high capacities in the North of the Biosphere Reserve reveal the location of the B1 Wiener Straße. All the capacity maps show the capacities of landscape elements within the open land of the Wienerwald Biosphere Reserve. Also, capacities of the landscape services Transportation are only assessed when in the open land, which explains the interruptions within the visualisation of the roads.

Landscape elements: High scored classes within the capacity matrix are paved open areas including parking areas and storage sites (4), public transport (5), roads primary (5), roads secondary (5), roads tertiary (4).

Three clusters with highest INDEC: Cluster 1 (INDEC 374,63) refers to an area at the B210 Badner Straße between the Helenental and Schwechatbach. Dominated by the primary road in size the polygon connects recreational trails, public transport, secondary and residential roads and areas for parking. Cluster 2 (INDEC 227,43) is located in Alland at the connection of A21 and B11. Cluster 3 (INDEC 167,04) refers to a section of the A21 connecting to the B11 close to Heiligenkreuz. This cluster again includes recreational trails, a parking area and a residential road.

Table 24: Three clusters with highest INDEC for Transportation. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes						
Cluster 1	374,63	126	1.188.937	RTR7	ROD3	PUT2	ROD6	AR59	AR60	ROD4
				72	28	12	9	2	2	1
Cluster 2	227,43	45	872.364	ROD3	RTR7	ROD4				
				33	9	3				
Cluster 3	167,04	44	588.371	RTR7	ROD3	AR60	ROD6			
				24	18	1	1			

5.2.17 Tourism facilities

Figure 27: Tourism facilities capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DEnsity Capacity (INDEC).

Spatial distribution: The landscape service Tourism facilities reveals a very fine pattern of capacity with small clusters in the whole area of the Biosphere Reserve. In the East high capacity clusters are located in Ottakring, Hütteldorf, Lainzer Tiergarten, Kalksburg (all Vienna). In the centre hotspots are visible in Purkersdorf, Pressbaum, Gablitz, Mauerbach, Kaltenleutgeben, and Krainerhütte and at the southern border of the Biosphere Reserve in Holzschlag and Nöstach.

Landscape elements: The classes with the highest scores in the capacity matrix are paved open areas including parking areas, storage sites (4), gastronomy (5), and accommodation (5).

Three clusters with highest INDEC: The first cluster (INDEC 28,34) is located in Purkersdorf including 21 landscape elements of recreational infrastructure, gastronomy, recreational trails and paved open areas used as parking spaces. The second cluster (INDEC 28,27) is located to Kaltenleutgeben close to the church. It emerges from recreational infrastructure (including a playground), recreational trails, a leisure and sport facility and a park. The third cluster (INDEC 25,47) is placed at the Steinhofgründe in Vienna and includes recreational infrastructure (playground Steinhofgründe), recreational trails, two parks or gardens and one accommodation.

Table 25: Three clusters with highest INDEC for Tourism facilities. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes			
Cluster 1	28,34	21	35.089	INF16	GST17	RTR7	AR60
				8	7	4	2
Cluster 2	28,27	23	25.186	INF16	RTR7	AR54	GD56
				18	3	1	1
Cluster 3	25,47	20	26.153	INF16	RTR7	GD56	ACC18
				9	8	2	1

5.2.18 Aesthetic information

Figure 28: Aesthetic information capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC)

Spatial distribution: The landscape service Aesthetic information shows low and medium capacities throughout the area of the Biosphere Reserve. Big clusters of very high capacity are visible in the North close to Klosterneuburg and Vienna/Nußdorf, in the South close to Groisbach and Baumgarten am Tullnerfeld in the NW.

Landscape elements: The highest scored classes within the capacity matrix are meandering, seminatural streams (5), seminatural, nutrient-poor lakes and pools (4), reed beds (4), nutrient-poor, extensive moist meadow (4), chasm woodland (5), abandoned fruit trees (5), nutrient-rich herbs (blooming aspect) (4), semidry managed meadows (4), nutrient-poor managed meadows and pastures (4), dry shrubland on limestone (5), thermophilous dry shrubland (5), semidry and dry grassland (5), dry forest edges (5), forest edges (5), *Pinus nigra* forests on dry grassland (5), species-rich slopes and margins (5), rock fragments and dry-stone walls (5), vineyard (5), hedges and shrubs (4), long-rotation woodland (4), hedgerows dominated by trees (5), tree rows and single trees (5), larch dominated meadows/pastures (5), riparian woodland (5), extensive orchards (5), cliffs (5), gardens or parks (5), greenery and planting (4), religious buildings (5), religious elements (5), artwork and monuments (5), castles and ruins (5).

Three clusters with highest INDEC: Cluster 1 (INDEC 515,68) is a large cluster (859 226m²) including 336 landscape elements located in the north of Vienna, covering the slopes of the Kahlenberg and Leopoldsberg from Grinzing to Nussdorf. Dominated by vineyards accompanied by parks or gardens and species poor field margins. In addition, it includes landscape structuring elements as forest edges, tree rows and single trees, tree dominated hedgerows and four extensive orchards. Cluster 2 (INDEC 243,55) is located close to Baumgarten am Tullnerfeld and includes vineyards, extensive orchards and cultivated fields as well as their fallow land, chasm woodland and managed meadows. Cluster 3 (INDEC 238,12) is located to Klosterneuburg (Buchberg, Kerblwiese) and integrates tree dominated hedgerows, vineyards, dry fallow land of fields and vineyards, parks or gardens, dry grassland and meadows, as well as 4 religious or artistic elements and a church.

Table 26: Three clusters with highest INDEC for Aesthetic information. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m²)	Distribution of LE classes											
Cluster 1	515,68	336	859.226	CU38	GD56	MG35	TR31	TR45	AR61	DR25	MD24	SB40	TR44	SB41	
				85	59	30	28	20	19	16	14	13	13	9	
				MD23	RC34	TR48	TH12	TR47	MG33	REL10	SB16	SB17	TH20		
				8	8	4	3	3	1	1	1	1	1		
Cluster 2	243,55	161	394.722	CU37	CU38	TR48	MD22	DR25	GD56	SB41	TR18	MD24	MD23	TR44	DR29
				19	19	19	16	14	11	11	10	7	6	6	4
				MG33	TR43	AR61	MG35	TR31	GD57	MD26	TH12	TH20	TR19	TR45	
				4	3	2	2	2	1	1	1	1	1	1	
Cluster 3	238,12	144	450.066	TR44	CU38	DR25	MD22	GD56	TR48	AR61	DR29	MD24	MD23	MG35	REL10
				31	28	15	14	13	5	4	4	4	3	3	3
				CU39	MG33	SB41	TR19	TR43	AR54	REL9	SB40	STA11	TH12	TR18	TR45
				2	2	2	2	2	1	1	1	1	1	1	1

5.2.19 Recreation

Figure 29: Recreation capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC)

Spatial distribution: For the LS Recreation capacities are shown throughout the area. Many high capacity areas are located in close proximity to settlements along the Biosphere Reserves North and East as Höflein an der Donau, Kritzendorf, Klosterneuburg and Vienna. In the South the hotspots are slightly more isolated with locations in Groisbach, Sankt Corona am Schöpfl and Obertriesting. In the Northwest high capacities are revealed along the border of the Biosphere Reserve from Sieghartskirchen and Freundorf until Wolfpassing.

Landscape elements: The classes with the highest scores according to the capacity matrix are meandering, seminatural streams (4), artificial standing water connected to groundwater (4), chasm woodland (5), abandoned fruit trees (5), semidry managed meadows (4), semi-dry and dry fallow land of fields and vineyards (4), semidry and dry grassland (5), dry forest edges (5), forest edges (5), *Pinus nigra* forests on dry grassland (4), vineyard (4), tree rows and single trees (4), larch dominated meadows/pastures (5), riparian woodland (4), extensive orchards (5), cliffs (4), abandoned stone quarries (4), abandoned gravel pits (4), lawns (4), cemeteries (4), religious buildings (4), museums (5), castles and ruins (4).

Three clusters with highest INDEC: Cluster 1 (INDEC 152,62) is located close to Wolfpassing at the Dopplerhütte along the slopes of the Eichberg (394m). 27 of the 102 landscape elements within this cluster are forest edges. Further 16 landscape elements are intermediate managed meadows and 10 hedgerows dominated by trees. Some gardens, dry meadows and extensive orchards are visible in the north and northwest of the cluster. Cluster 2 (INDEC 135,67) is located in the north of Vienna at the Kahlenberg. Vineyards, gardens or parks, forest edges and field margins dominate this cluster. Furthermore, it includes a view point and a recreational trail. Cluster 3 (INDEC 130,62) is located in the north of the Biosphere Reserve close to Höflein an der Donau and counts 87 landscape elements. Next to forest edges, extensive orchards, semidry meadows it includes gardens or parks, recreational trails, a cemetery, and altogether four religious buildings and elements.

Table 27: Three clusters with highest INDEC for Recreation. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m²)	Distribution of LE classes										
Cluster 1	152.,62	102	242.064	TR31	MD23	TR44	GD56	MD22	TR48	AR61	DR25	RTR7	CU38	MD13
				27	16	10	7	7	7	5	3	3	2	2
				SB17	TR18	TR43	MG33	MG35	MG36	REL10	TR19	TR45	TR47	
				2	2	2	1	1	1	1	1	1	1	
Cluster 2	135,67	91	213.594	CU38	GD56	TR31	MG35	AR61	DR25	INF16	RC34	TR45		
				21	18	16	11	7	7	2	2	2		
				TR48	AR59	MD23	MG33	MG36	RTR7	SB17	VPS15	WT5		
				2	1	1	1	1	1	1	1	1		
Cluster 3	130,62	87	208.608	TR31	TR48	MD22	GD56	RTR7	DR25	TR18	TR44	AR61	DR29	MD26
				14	14	12	6	6	4	4	4	3	3	3
				REL10	REL9	TR19	AR59	CU38	GD58	MD23	SB17	TR43	TR45	TR47
				2	2	2	1	1	1	1	1	1	1	1

5.2.20 Cultural and artistic information

Figure 30: Cultural and artistic information capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENSITY Capacity (INDEC)

Spatial distribution: High capacities of the LS Cultural and artistic information are located in the north of the Wienerwald close to the Danube River. Starting in the North of the city Vienna up to Klosterneuburg, Kritzendorf and Höflein an der Donau. Furthermore, isolated spots of high capacity are visible in Ottakring and Mauer (Vienna) and in the South Heiligenkreuz, Groisbach, Burg und Graben, Reitel-Graben. In the NW the area close to Baumgarten am Tullnerfeld up until Königstetten reveals spots of high capacity.

Landscape elements: High scored classes of the service artistic and cultural information within the capacity matrix are dry shrubland on limestone (4), rock fragments and dry-stone walls (5), vineyard (5), tree rows and single trees (5), larch dominated meadows/pastures (5), larch dominated meadows/pastures (5), extensive orchards (5), gardens or parks (5), cemeteries (5), isolated buildings (4), trails educational (4), religious buildings (5), artwork and monuments (5), archaeological sites (4), museums (5), castles and ruins (4).

Three clusters with highest INDEC: Cluster 1 (INDEC 106,6) is located in Vienna's 23rd District, close to the Lainzer Tiergarten. Nestled in the transition area from forest to the settlements of Vienna, it is dominated by tree rows and single trees, vineyards and field margins as well as gardens, parks and isolated buildings. Cluster 2 (INDEC 104,1) refers to an area in Vienna's 19th district/Neustift am Walde. It agglomerates mainly of gardens or parks and vineyards together with dry fallow land of vineyards, tree rows, single trees and isolated buildings. The third cluster's (INDEC 100,59) location is in Klosterneuburg Weidling. It is an aggregation of small and medium size polygons of 19 different classes, mainly vineyards, some gardens or parks of isolated buildings. Furthermore, four religious elements are contributing to the cluster.

Table 28: Three clusters with highest INDEC for Cultural and artistic information. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes												
Cluster 1	106,6	73	160.676	TR45	GD56	CU38	MG35	AR61	MG33	DR30	TR48	WT4	DR25	REL10	REL9	TR19
				16	10	9	8	7	6	5	5	3	1	1	1	1
Cluster 2	104,1	70	163.047	GD56	CU38	AR61	DR25	TR45	MD24	MG35	MG33	TR44	DR29	MD23	REL10	WT6
				19	15	7	7	7	4	3	2	2	1	1	1	1
Cluster 3	100,59	66	165.407	CU38	GD56	AR61	REL10	TR44	DR25	MG35	SB41	TR45	CU37	MD24		
				17	8	6	4	4	3	3	3	3	2	2		
				TR19	TR43	TR48	AR54	CAS14	DR29	MD22	WT3					
				2	2	2	1	1	1	1	1					

5.2.21 *Spiritual and historic information*

Figure 31: Spiritual and historic information capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DENsity Capacity (INDEC)

Spatial distribution: The LS spiritual and historic information shows high capacity in the north of the Wienerwald Biosphere Reserve, starting from Baumgarten am Tullnerfeld to Kleinstaasdorf, Höflein an der Donau, Kritzendorf and Klosterneuburg. Also, the northern area of Vienna shows dens clusters of high capacity. Additionally, isolated spots are visible in Mauer an area in the 23rd district of Vienna as well as in the in the south (Heiligenkreuz, Groisbach, Holzschlag, Berg und Graben, close to the ruin of Sankt Pankraz and Klauswies) and west of the Biosphere Reserve (Wöllersdorf, Innerfurth and Ödengraben).

Landscape elements: The classes with the highest capacity according to the capacity matrix are straight, seminatural streams (4), meandering, seminatural streams (4), chasm woodland (5), chasm woodland (5), dry shrubland on limestone (4), rock fragments and dry-stone walls (5), vineyard (5), tree rows and single trees (5), larch dominated meadows/pastures (5), extensive orchards (5), gardens or parks (4), cemeteries (5), isolated buildings (4); religious buildings (5), religious elements (4), artwork and monuments (4), archaeological sites (5), museums (5), castles and ruins (5).

Three clusters with highest INDEC: The cluster with the highest INDEC (96,01) is located in Klosterneuburg and is an agglomeration of big to small polygons. It is dominated by vineyards, gardens or parks of isolated buildings and includes four religious elements (e.g. the “Gabelkreuz Weidling” and a ruin. The second ranked cluster (INDEC 94,8) is located in Vienna, in Obersievering. Emerging from 64 landscape elements, it contains mainly gardens or parks (18 landscape elements), vineyards (15 landscape elements). The third cluster (INDEC 89,79) is also in Vienna at the Lainzer Tiergarten. It includes mainly tree rows and single trees, vineyards and gardens or parks of isolated buildings.

Table 29: Three clusters with highest INDEC for Spiritual and historic information. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m²)	Distribution of LE classes											
Cluster 1	96,01	62	162.631	CU38	GD56	AR61	REL10	TR44	DR25	SB41	TR45	CU37	MD24	TR19	
				17	8	6	4	4	3	3	3	2	2	2	
				TR43	TR48	CAS14	DR29	MD22	WT3						
				2	2	1	1	1	1						
Cluster 2	94,8	64	147.271	GD56	CU38	AR61	TR45	DR25	MD24	TR44	DR29	MD23	MG33	REL10	WT6
				18	15	7	7	6	4	2	1	1	1	1	1
Cluster 3	89,79	60	142.428	TR45	CU38	AR61	GD56	MG33	TR48	DR30	WT4	DR25	REL10	REL9	TR19
				16	8	7	7	6	5	4	3	1	1	1	1

5.2.22 Science and education

Figure 32: Science and education capacity in the Wienerwald Biosphere Reserve. The colour ramp distinguishes clusters of different INTensity and DEnsity Capacity (INDEC)

Spatial distribution: The high capacities of the LS Science and education are concentrated to the north and south to southwest of the Wienerwald Biosphere Reserve. In the east and northeast, the high capacity areas are bordering Vienna. Especially the Leopoldsberg and Kahlenberg region in the north, as well as Hietzing and the Maurer Wald in the northeast, show areas of high capacity for science and education. In the south the map shows agglomerations of high capacity areas close to Altenmarkt an der Triesting, Burg und Graben and Koissbach between the elevations of Peilstein und Kuhberg. Continuing along the border of the Wienerwald Biosphere Reserve in the south, clusters with high capacity are visible around Wöllersdorf and Innermanzing in the southwest. High capacity clusters in the north and northwest are located in Sieghartskirchen and in proximity to Freundorf, Tulbing, and Wolfpassing and again in Höflein an der Donau, Kritzendorf and Klosterneuburg.

Landscape elements: Classes with the highest scores according to the capacity matrix are meandering, seminatural streams (5), straight, seminatural streams (4), seminatural, nutrient-poor lakes and pools (4), nutrient-poor, extensive moist meadow (4), chasm woodland (4), abandoned fruit trees (4), dry shrubland on limestone (4), thermophilous dry shrubland (4), semidry and dry grassland (4), dry forest edge (5), forest edge(5), species-rich slopes and margins (4), rock fragments and dry-stone walls (5), vineyard (4), hedges and shrubs (4), hedgerows dominated by trees (4), tree rows and single trees (5), larch dominated meadows/pastures (5), riparian woodland (4), extensive orchards (5), cliffs (4), trails educational (4), religious buildings (4), archaeological sites (5), museums (5), castles and ruins (5).

Three clusters with highest INDEC: The first cluster (INDEC 183,15) is located in the north of the Viennese part of the Wienerwald Biosphere Reserve in Döbling and Nußdorf on the slopes of the Kahlenberg. It emerges from a mosaic of polygons of 123 different elements. It includes mainly gardens or parks of the settlements and isolated buildings, vineyards and fallow land, forest edges, tree dominated hedgerows, hedges and shrubs and some the riparian woodland accompanying the Schreiberbach. The second cluster (INDEC 159,52) is close to Groisbach in the south of the Biosphere Reserve mainly consisting of tree rows and single trees as well as tree dominated hedgerows. It includes managed meadows ranging from semidry meadows, pastures and cultivated fields in the north and middle of the cluster to moist meadows in the south of the cluster. In the east of the cluster in proximity of the settlements a few extensive orchards accompanied by tree rows and single trees are visible. The third cluster (INDEC 159,21) is in the north of the Biosphere Reserve close to the Dopplerhütte at the slopes of the Eichberg in Königsstetten. Framed by forest edges, it includes a wider range of landscape elements, like different managed meadows, tree related classes (as tree dominated hedgerows, tree rows, and extensive orchards) and isolated buildings.

Table 30: Three clusters with highest INDEC for Science and education. The INDEC value, the frequency of landscape elements (N LE), the area of the clusters and the distribution of landscape elements per class are reported. For the LE classes' code, refer to Table A 3 and Table A 4.

CLUSTER	INDEC (adim)	N LE	Area (m ²)	Distribution of LE classes											
Cluster 1	183,15	123	287.609	GD56	CU38	TR31	AR61	MD24	TR44	DR25	RC34	TR45	SB41		
				33	28	13	7	7	7	6	6	5	2		
				TR47	TR48	MD23	MG33	SB17	SB40	TH21					
				2	2	1	1	1	1	1					
Cluster 2	159,52	106	255.918	TR45	TR44	MD22	MD24	TR31	CU37	MD13	MD23	TR48	AR61	DR29	
				38	19	10	5	5	4	4	4	4	3	3	
				DR25	MG33	MD26	SB41	TR47							
				2	2	1	1	1							
Cluster 3	159,21	108	244.902	TR31	MD23	TR44	GD56	TR48	AR61	MD22	TH12	DR25	SB41	CU38	
				28	15	11	7	7	6	6	6	3	3	2	
				MD13	SB17	TR18	TR43	MG33	MG36	REL10	SB16	TR19	TR47		
				2	2	2	2	1	1	1	1	1	1		

5.3 Selected landscape services for a BRIM

After analysing the capacity maps and related INDEC, we could select the landscape services for a BRIM. As explained in the section 4.10, combining the selected landscape services for the BRIM merges the qualities of representativeness, communication and relevance for the Biosphere Reserve Management. The representativeness is provided by landscape services addressed by a high amount of landscape services classes and therefore high capacity within the Wienerwald Biosphere Reserve. Landscape services that convey clarity and are easily understandable for the Wienerwald Biosphere Reserve Management or potentially involved stakeholders produce a set of indicators useful also for communication purposes. Additionally, we emphasised landscape services with an overarching character, but we also considered a balance between ecological and socio-cultural aspects, as both are very relevant for the Wienerwald Biosphere Reserve Management. The combination of the different aspects can be depicted as a flower, where each petal represents a different sphere of the Biosphere Reserve environment (Figure 33). Following the flower scheme, the eight selected landscape services are embedded into a BRIM flower, which highlights how each service corresponds or partially overlaps to a sphere. In this way, an integral monitoring of the Wienerwald Biosphere Reserve can be guaranteed (Figure 34).

Figure 33: Scheme showing the spheres of the BR environment.

Figure 34: Landscape services for a BRIM for the Wienerwald Biosphere Reserve.

Local climate and air regulation is related to the Atmosphere, it is an overarching service, and it is important due to closeness of highly populated areas, such as Vienna, that is going to be subjected to more and heat-related problems, due to the climate change effects. Refugium is related to Biosphere and it is a very relevant indicator or proxy for conservation value. The service Soil retention belongs to Lithosphere, it holds a strong communication potential, as the concept of soil erosion can be easier to grasp than the similar service Soil formation. The service Water supply belongs to the Hydrosphere, it is very easy to communicate and to understand. The **Information** services Recreation, Aesthetics, Science and Education belong to the Anthroposphere. Science and Education is a crucial indicator for every biosphere reserve, providing this is one of the main tasks for Biosphere Reserve management, in accordance to the “Sevilla Strategy” goal III. Recreation has a pivotal importance due to closeness of the reserve to Vienna. Moreover, this indicator is also included in the “Sevilla Strategy”. Both Aesthetic information and Recreation have a strong focus on open land elements as well as on additional spatial indicators. Last, the service Food is obviously a fundamental indicator of performance of the Biosphere Reserve. Since this service includes both wild and cultivated goods, it is more inclusive than the service Cultivation. Food belongs to more than one sphere, being part of the Anthroposphere, due to its connection to human basic needs, but also part of the Biosphere, because of the wild products.

6 Discussion

Throughout the project's meetings and developments both criticism and praise were raised and discussed by the project team, colleagues and project partners. The main topics are summarized in the following sub-chapters.

6.1 Integrated discussion on the landscape services capacity in the Biosphere Reserve

The capacity maps showed a great variability in the provision of service capacity. Some services were highly represented in the Biosphere Reserve (e.g. Soil formation and Nutrient Regulation), while others were underrepresented (e.g. Raw Materials). **Regulating** services showed very similar patterns across the reserve, and were related mainly to seminatural open land classes, whereas some **Information** services and some **Carrier** services relied mainly on additional spatial indicator classes. The cultivated large area in the NW of the Biosphere Reserve was hardly targeted by the **Regulating** services, while high INDEC values for the **Provisioning** service Food and for the **Carrier** Cultivation were revealed.

Even within the service categories, the distribution of the clusters in the Biosphere Reserve was not always coupled, as in the case of provisioning services. This fact shows that the service categorization might be not the most straightforward way to synthesize services, as done in previous studies (see for instance Hermann et al. 2014), as this might lead of to excessive reduction of variability and information that each service can deliver.

Another interesting point is the scale discrepancy among services. Services can have scale mismatches and delocalized causes and effects. For instance, the **Regulating** service climate regulation has a much higher effect which exceeds the single service providing unit "landscape element". On the other hand, there are services with a very punctual influence on the landscape. To take this aspect fully into account, landscape service capacities might be weighted differently (Burkhard et al. 2012).

Regulating services showed very similar patterns across the reserve, with larger clusters mostly concentrated in the northern region of the Biosphere Reserve, and other high INDEC values detected in the most northern corner of the Biosphere Reserve, in the proximity of Höflein an der Donau. Other high capacity areas were often detected between Klosterneuburg and the municipality of Vienna. Other important concentrations of clusters are generally located in the southern sector, NE of Altenmarkt an der Triesting, and in the proximity of Laaben, in the very western sector of the reserve. The cultivated large area in the NW of the Biosphere Reserve was hardly targeted. Although the distribution of the clusters in the Biosphere Reserve was similar, some **Regulating** services showed much higher capacity than others. Soil formation, soil retention, pollination and nutrient regulation showed much higher INDEC values and much larger and dense clusters than the other services. This fact highlights that **Regulating** services might be related to higher levels of naturalness.

Similarly to **Regulating** services, **Habitat** services are also related to the level of naturalness; this is more valid for Refugium than for Nursery. Indeed, the Biosphere Reserve includes very few natural landscape elements, as most of them can be considered semi-natural (e.g. streams, managed meadows, forest edges etc.). Moreover, most landscape elements with

high naturalness level are generally rare and small in size, and for this reason they couldn't be always highlighted through the method employed. Nursery and Refugium show similar distribution patterns in the Biosphere Reserve, with Refugium being very well represented and associated to very high INDEC values. Their service capacity is generally very high and very spread in the reserve, and more concentrated in the N of the Biosphere Reserve. Other significant areas can be found in the southern part of the Biosphere Reserve, in the very large mosaic of meadows and fields just N of Altenmarkt an der Triesting. Most of the open land classes can potentially provide Refugium service, depending on their level of human disturbance or management.

Provisioning services include quite different services, from Food to Medicinal Resources to Raw Materials and Genetic Resources. Due to this heterogeneity, no congruent pattern in the service distribution can be detected. On the one hand, Food shows high INDEC values and large clusters, associated to the NW cultivated sector of the Biosphere Reserve; on the other hand, the services Raw materials and Genetic Resources have negligible service capacity in the Biosphere Reserve. One reason behind this variability lies in the definition of Raw Resources, which, according to BRIMSEN approach, comprise sand and gravel. The Biosphere Reserve does not include many active quarries. As to Genetic Resources, they mainly refer to forestry, which is excluded from the study area.

The category of **Carrier** services relates to very different classes for each landscape service. Whereas the service Cultivation only refers to classes of the open land, mainly large sized agricultural structures (cultivated fields etc.) displayed in the north and northwest, the Tourism facilities and Transportation are strongly depended on classes of artificial landscape elements provided both by the mapping of the open land as well as additional spatial indicators.

The service Tourism facilities shows a fine display in capacities throughout the Biosphere Reserve. Since the classes of gastronomy and accommodation have received the highest scoring for Tourism facilities in the capacity matrix, this service could benefit from the inclusion of the settlements to the analysed data (service Tourism facilities max. INDEC = 28,34 compared to e.g. landscape service Recreation max. INDEC = 152,62).

The service Transportation strongly visualises the main traffic connections within the Wienerwald Biosphere Reserve. The analysis of the clusters showed that the biggest cluster was based on a street, but also includes recreational trails, public transport, smaller roads and areas for parking. Therefore, high capacity areas of Transportation seem to represent both open land strongly influenced by motorized traffic but also highly connected to human accessibility of the area (Abraham et al. 2009). Since the water-related habitats of Wienerwald are too small not deliver a Transportation service, we did not consider them as such. However, in other regions watercourses might provide the **Carrier** service Transportation (e.g. Danube River), and this is true not only for running waters but also for standing waters.

The category of **Information** services shows hotspots close to settlements throughout all the services, mainly along the border of the Biosphere Reserve. High capacities are always visible in the north of the Biosphere Reserve. The centre and west of the Biosphere Reserve show lower capacity in all **Information** services, compared to the rest of the Biosphere Reserve. However, differences are visible in the number of classes with capacity of each service. The **Information** services strongly refer to both seminatural open land classes as well as transformed elements of the additional spatial indicators. Compared to other studies (see

Hermann et al. 2014) the detailed classification of landscape elements within a joint approach for both disciplines (eco and sociocultural) allowed for a differentiated consideration of seminatural elements within the **Information** services. Thereby the possibility of education and appreciation by experiencing nature could be assessed (Hughes et al. 2018; Prévot et al. 2018; Dean et al. 2019). The concentration of areas of high capacity of most **Information** services close to settlements can be linked to the transformed landscape elements. The inclusion of those connects nature to both its (socio-) cultural perception (Weichhart 2006) and accessibility (Reimoser et al. 2008; Bell et al. 2007). The well spread pattern of the service Aesthetic information throughout the Wienerwald Biosphere reserve can be linked to the fragmented open land landscape of the Wienerwald.

6.2 Considerations on data availability and data treatment

6.2.1 *Data sources*

In our study we aimed to address a broad range of landscape services to show the diversity of benefits the landscape provides for humans. As faced in other projects (Wrbka et al. 2012) the availability and compatibility of geodata was a main issue raised during the work on the project.

Especially the assessment of the **Carrier** and **Information** services categories strongly depend on the availability of data additional to the mapping of the open land habitat types. Therefore, it was necessary to find a way to balance data varying in sources, age and level of accuracy (see section 4.8). Nevertheless, the dismissal of landscape services was required for those cases where the provision of sufficient data was not accomplishable.

The project team faced both decisions regarding inclusion of data based on availability as well as exclusion of data based on accuracy or quality of results.

Therefore, we excluded the data of the settlements because its very broad categorisation was incomparable to the detailed quality of the open land habitat types. Second, we dismissed parts of the data in the southeast due to a great heterogeneity in the level of accuracy.

The inclusion of data of the forest was limited to the forest edges. The advantages of focussing on the open land exclusively are the differentiated results produced. However, the effect of including the forest can roughly be visualised by comparing the two maps of Figure 35, created using different methods. The disproportional capacity performance of the forest (Figure 35a) overlays the performance of the open land (Figure 35b). Therefore, by excluding the forest meaningful results on the assessment of the landscape service capacities of the open land of the Wienerwald can be achieved.

Figure 35: Comparison between the inclusion and exclusion of the forest.

6.2.2 Data classification

Another critical aspect was the classification of the data. The data of the open land is based on a mapping of the open land habitat types from an ecological point of view. On these grounds the open land habitat types referred to very detailed categorisation within the natural or semi-natural elements, whereas the other more anthropogenic classes provide broad information only. Since the further revision and amendment of the data was not possible, these broad classes (e.g. gardens or parks (GD56) or unvegetated open areas with compacted soil including parking areas and storage sites (AR59)) represent an imprecision in the evaluation progress as well as in the results. Other classes, like settlements, were excluded from the data selection.

6.3 Reflections on the methodology

6.3.1 *The capacity matrix: advantages and disadvantages*

Capacity matrices have proved to be a valuable method for the assessment and evaluation of landscape services (Hermann et al. 2014; Burkhard et al. 2012; Campagne et al. 2018). They provide a flexible, integrative and time and cost-efficient approach to collect and integrate expert knowledge as a source of information (Campagne et al. 2018; Jacobs et al. 2015). The attractiveness of the matrix approach results from its flexibility concerning level of detail and level of abstraction from rather simple to highly complex. Its potential to integrate all kinds of data, from expert-scores to statistics, interview data, measurements or high-end model outcomes makes it applicable in data-poor as well as data-rich environments. Results based on the flexible 0-5 ranking system and the linkage to geo-biophysical spatial units in ecosystem service maps provide wide application ranges in science and in decision making (Burkhard et al. 2014). On the other hand, there are several uncertainties related to the matrix method applied for landscape analyses Hou et al. (2013), which we could experience in BRIMSEN as well.

In BRIMSEN the expert knowledge emerged from the teams cooperating in the project. Therefore, the consensus approach provided productive input and discussion, minimizing ambiguities in the definitions and improving the development of the project. Furthermore, as also pointed out by Jacobs et al. (2015) the balancing effect of the consensus method allowed to progress on decisions where uncertainty (due to lack of data or knowledge) might have been blocking the individual. Nevertheless, an individual scoring beforehand to support the possibility of statistical analysis could be considered. Moreover, expanding the number of experts involved and thus optimizing scientific credibility shall also improve the method (Campagne et al. 2018; Jacobs et al. 2015). In fact, there is a high dependence on the observer's experience, knowledge and objectivity which services are supposed to be relevant and how to value them (Burkhard et al. 2012).

Although the issue of double counting of elements was avoided by agreement on distinct definitions or a splitting on values within strongly connected services (Wrbka et al. 2012), the distinction between the assessment of a general capacity of classes/elements while strongly referring to local/regional conditions and peculiarities of the Wienerwald Biosphere Reserve (also due to the Wienerwald explicit data of the open land habitat types) revealed contextual inconsistencies. The possibility of a direct transferability of the scores to other biosphere reserves or nature protection areas should be object of critical revision. Another limitation of the capacity matrix and of the study was that the actual condition of the landscape element and the influence of seasonal aspects were not considered addressed. In this sense the capacity score assigned wasn't adjusted with a qualifier factor, as done in other studies (e.g. Hermann et al. 2014; Hainz-Renetzeder et al. 2014).

In line with Campagne et al. (2018) the focus on the assessment of capacities does not allow for the consideration of trade-offs or neighbouring effects (positive or negative) within the matrix, therefore the complexity and multifunctionality of ecosystems is embodied insufficiently.

Finally, although careful preparation of the materials to assure transparency and time efficiency was aimed for within the BRIMSEN capacity matrix workshop, the goal of scoring all 80 classes within one day in consensus could not be achieved. For which reason we see an accessible size of the matrix as well as and realistic time management throughout the scoring as key issues to avoid revisions, in the achievement of a consistent result.

Based on this experience we are aware that no final solution for highly complex ecosystem and landscape service assessments has been found yet and that related challenges are still manifold.

6.3.2 *Intensity and Density of service Capacity: advantages and disadvantages*

The INDEC is a newly proposed methodology that allowed us to overcome several issues related to data quality. In fact, through the INDEC application all landscape elements were treated at the same level, being all transformed into point data. The second advantage of the method is that it conveys multiple simultaneous information of density and of cumulative intensity, based on capacity score and the size of service capacity. The approach had also the great advantage of addressing very different services ranging from ecological to socio-cultural aspects. Indeed, the INDEC was very adequate for the combination of different data sources, which, otherwise, would have to be merged and modified in a GIS environment. Combining and merging data from different sources means having to deal with different resolution, age, spatial reference, different metadata and so on. The INDEC proved very efficient and successful in overcoming the data discrepancies and therefore it could be particularly useful in projects with limited financial resources. Thanks to the INDEC, there was no modification in the original data sets of open land data and in the additional spatial indicators. Moreover, our methodology highlighted the spatial connectivity of landscape elements, thanks to the creation of clusters equally dependent on the intensity of the capacity, assigned through the capacity matrix, and on the density of the landscape elements, i.e. their proximity. Therefore, the method allowed an appropriate balance among isolated and large landscape elements (e.g. cultivated field, meadow) and small-scale but numerous landscape elements, such as forest edges, tree rows and benches. By adopting a method that applies a simple area-weighted factor to each polygon, most of the small-scale elements would be overlooked.

The INDEC is based on a tailored landscape services assessment. Nevertheless, the methodology could be efficiently transferred to other data sets, in any other region, provided that the capacity matrix scores are revised. The efficiency of the INDEC makes it fruitful for applied analyses, such scenarios development. During the project duration, some exercises on scenario development were carried out. For instance, we hypothesized the effects of management measures on the future provision of landscape services. Realistic measures might be extensive woodlots logging, change in the management of meadows, conversion of cultivated fields into vineyards or another cultivation, and so on. The INDEC might quite rapidly provide projections on the change of landscape service provision, based on each of such measures. Similarly, the method could be employed as monitoring tool, by simply revising the matrix scores and the data sets. Last but not least, the INDEC is an approach fitting with the rationale behind the capacity matrix, since the size of landscape elements areas is not explicitly taken into account.

Shifting to the critical issues encountered in the development of the INDEC, we noticed that in some cases the density was a dominant factor over the intensity. This applies to the forest edges, which were so abundant they were often included in the clusters of highest capacity. On the other hand, this method relies on the quality of the data set: in BRIMSEN, we dealt with an “unbalanced” dataset, with high number of classes with few landscape elements (e.g. thermophilous dry shrubland with two landscape elements), and a low number of classes with many landscape elements: forest edges (11032 landscape elements); trails (7823); intermediate managed meadows (3283); hedgerows dominated by trees (3041). A critical aspect in the development of the methodology was “translating” the difference in

landscape services capacities into a buffer. In other words, what is the “spatial” proportion between a score “2” and a “5”? Another technical issue regarded the transformation into points: depending on the shape of the polygon, the centroids of polygons sometimes were outside the area of the polygon they belonged. This aspect remained unresolved.

The INDEC and the method employed in BRIMSEN in general might be considered rather simplistic in the sense that they do not address disservices, i.e. conditions blocking the capacity of certain services, nor neighbouring aspects, i.e. influence of other landscape elements on the scoring for the service. Moreover, the INDEC is static: no temporal dynamism was considered.

As a secondary criticism, it is worth mentioning that the huge data set handled needs powerful hardware and software. However, this aspect is not limited to the INDEC.

6.4 Management implication

BRIMSEN provided very useful tools for a more effective management of the Wienerwald Biosphere Reserve. By focussing on the open land areas of the Biosphere Reserve, high value to the small-scale elements was provided. In fact, as shown in Figure 35, the disproportional extent and relevance of the forest generally suffocates the role of the open land. Therefore, addressing the open land only was very useful for highlighting its multifunctionality. The landscape service maps can be fruitfully adopted by the management staff of the Reserve to identify hotspot for certain service, and to orient their measures consequently. For instance, currently farmers receive financial support based on the size of their land, and not based on the number of valuable small-scale elements included. Stressing the role of the overlooked open land elements might change their attitude towards the conservation of their ecological integrity.

BRIMSEN proposed a set of indicators for a BRIM, which can be employed as indicators for monitoring different aspects for the reserve. The variegated set of the indicators opens an integrated vision for the Reserve, where each service contributes to enhance its sustainability performance. According to von Geisler et al. (2010) the differentiation between sustainable and non-sustainable development requires the analysis of the interactions between indicators within a socio-ecological system.

Through BRIMSEN we have concretely contributed the Lima Action Plan 2016-2025 (UNESCO 2016), in particular to the Outcome A7 “Biosphere Reserves recognized as sources of ecosystem services”, actions A7.1 “Identify ecosystem services and facilitate their long-term provision, including those contributing to health and wellbeing” and partially A7.3 “Implement programmes to preserve, maintain and promote species and varieties of economic and/or cultural value and that underpin the provision of ecosystem services”. In fact, the project allowed a tangible testing of the landscape services indicators, thus enhancing the awareness of the multifunctional/multi-service role of the Biosphere Reserve. BRIMSEN gave a contribution to the fulfilling of knowledge gaps related to natural capital, and enhancing awareness of values related to natural resources (see section 2.1). This contribution proved useful for the full realization of the biosphere reserve as model for Sustainable Development (Ruoss 2013).

BRIMSEN contributed also to the implementation of the socio-ecological monitoring system named “Sustainability Geoscope” (Lucht et al. 2004; Lotze-Campen et al. 2008). The Geoscope vision aims at an instrument for systematic collection and analysis of compatible natural-scientific and socio-economic data that enable a validation of integrated views of society-nature dynamics.

7 Conclusions

BRIMSEN provided an innovative tool named INDEC to simultaneously assess the whole range of landscape services. 22 capacity maps were created and a set of eight indicators for a BRIM was proposed in this study. Although the assessment performed through the capacity matrix has a local character, the tool might be transferable to other reserves and other regions.

Several aspects might be improved and further tested in the next studies. The most interesting one would be to analyse the spatial interaction among landscape services, in order to answer stimulating research questions like: can we see trade-offs when overlapping the selected services?

In a second step, one could explore positive and negative interactions among services. Some proposed interactions might be the following:

- Recreation vs Local climate regulation. We expect a positive interaction, with an enhancement in capacity for areas accessible and close to water-related open land elements or/and sheltered by trees.
- Aesthetics vs Refugium. We expect a positive interaction, with an enhancement in service capacity for mosaic areas (density of small elements) with reasonable or high level of naturalness.
- Food vs Local climate regulation. We expect a low overlapping of spots with high capacity. Areas with high values for food and low values for climate regulations should be managed in a way to enhance the local climate capacity (e.g. enhancement of diversity of landscape elements).

Moreover, as done by Hermann et al. (2014), the services could be eventually synthesized through a spider web diagram, to obtain an overview on the main trends per service category.

Since BRIMSEN focuses on service capacity only, no information on service demand is currently available. A future project might focus on this aspect, finding gaps between supply and demand (Burkhard et al. 2012).

Another stimulating project might include the development of the BRIMSEN concept and INDEC for the whole area of the Wienerwald Biosphere Reserve, by including forest and settlements. The results would be comparable to those presented in this study. In addition to this, since the INDEC method was tailored for a regional approach, it would be fascinating to tune it at the local scale, by selecting a set of municipalities within the Reserve. In this way, communication and inclusion of stakeholders or the public might be desirable.

As mentioned in the section 6.3.2, the methodology could be further tested for scenario development for management. Since the concept of landscape services is not mainstreamed yet, a fruitful idea could be the creation of a dedicated app to help the general public understand it better.

It has been suggested that we have now entered the Anthropocene, an epoch in which we have seen a 'quantitative shift in the relationship between humans and the global environment' (Steffen et al. 2011). If we are to solve the problems posed by global environmental change it has been argued that we need better coordinated international

research that places equal emphasis on the social and natural sciences (Perrings et al. 2011). A focus on ecosystem and landscape services is potentially one way in which a balanced perspective might be achieved (Potschin et al. 2011b).

8 References

- Abraham, A., Sommerhalder, K., & Abel, T. 2009. Landscape and well-being: A scoping study on the health-promoting impact of outdoor environments. *International Journal of Public Health* 55: p.59–69.
- Arnberger, A. et al. 2002. *Machbarkeitsstudie Wienerwald - Eignung des Wienerwaldes für einen Nationalpark oder Biosphärenpark*.
- Banos-González, I., Martínez-Fernández, J., & Esteve-Selma, M.A. 2016. Using dynamic sustainability indicators to assess environmental policy measures in Biosphere Reserves. *Ecological Indicators* 67: p.565–576. Available at: <http://dx.doi.org/10.1016/j.ecolind.2016.03.021>.
- Baral, H., Keenan, R.J., Fox, J.C., Stork, N.E., & Kasel, S. 2013. Spatial assessment of ecosystem goods and services in complex production landscapes: A case study from south-eastern Australia. *Ecological Complexity* 13: p.35–45.
- Bastian, O., Grunewald, K., Syrbe, R.U., Walz, U., & Wende, W. 2014. Landscape services: the concept and its practical relevance. *Landscape Ecology*.
- Bastian, O., Haase, D., & Grunewald, K. 2012. Ecosystem properties, potentials and services - The EPPS conceptual framework and an urban application example. *Ecological Indicators* 21: p.7–16. Available at: <http://dx.doi.org/10.1016/j.ecolind.2011.03.014>.
- Bastian, O., & Schreiber, K.-F. 1999. *Analyse und ökologische Bewertung der Landschaft* 2nd ed. Heidelberg, Berlin: Spektrum Akademischer Verlag Heidelberg.
- Bell, S., Tyrväinen, L., Sievänen, T., Pröbstl, U., & Simpson, M. 2007. Outdoor Recreation and Nature Tourism: A European Perspective. *Living Reviews in Landscape Research* 1.
- Bianchin, S. et al. 2012. *TransEcoNet Ecosystem services - Implementation of the concept of ecosystem services in protected and non-protected tranboundary regions WP5 - Output XV*. Wien.
- Biosphärenpark Wienerwald Management GmbH. Lebensregion Biosphärenpark Wienerwald - Region. Available at: <https://www.bpww.at/de/themenseiten/region> [Accessed April 23, 2019a].
- Biosphärenpark Wienerwald Management GmbH. Lebensregion Biosphärenpark Wienerwald - Zonierung. Available at: <https://www.bpww.at/de/themenseiten/zonierung> [Accessed April 24, 2019b].
- Brenner, H., Drozdowski, I., & Mrkvicka, A. 2015. *Wälder im Wienerwald*. Biosphärenpark Wienerwald Management GmbH.
- Brown, C., Burns, A., & Arnell, A. 2018. A Conceptual Framework for Integrated Ecosystem Assessment. *One Ecosystem* 3.
- Burkhard, B., Kandziora, M., Hou, Y., & Müller, F. 2014. Ecosystem service potentials, flows and demands-concepts for spatial localisation, indication and quantification. *Landscape Online* 34(1): p.1–32.
- Burkhard, B., Kroll, F., Müller, F., & Windhorst, W. 2009. Landscapes' capacities to provide ecosystem services - A concept for land-cover based assessments. *Landscape Online* 15(1): p.1–22.
- Burkhard, B., Kroll, F., Nedkov, S., & Müller, F. 2012. *Mapping ecosystem service supply, demand and*

- budgets*. Elsevier Ltd. Available at: <http://dx.doi.org/10.1016/j.ecolind.2011.06.019>.
- Burkhard, B., & Maes, J. eds. 2017. *Mapping Ecosystem Services*. Sofia: Pensoft Publishers.
- Campagne, C.S., & Roche, P. 2018. May the matrix be with you! Guidelines for the application of expert-based matrix approach for ecosystem services assessment and mapping. *One Ecosystem* 3: p.e24134. Available at: <https://oneecosystem.pensoft.net/articles.php?id=24134>.
- Carr, M.H. et al. 2017. The central importance of ecological spatial connectivity to effective coastal marine protected areas and to meeting the challenges of climate change in the marine environment. *Aquatic Conservation: Marine and Freshwater Ecosystems* 27(January): p.6–29.
- Chee, Y.E. 2004. An ecological perspective on the valuation of ecosystem services. *Biological Conservation* 120(4): p.549–565.
- Clerici, N., Paracchini, M.-L., & Maes, J. 2014. *Land-cover change dynamics and insights into ecosystem services in European stream riparian zones*.
- Crossman, N.D. et al. 2013. A blueprint for mapping and modelling ecosystem services. *Ecosystem Services* 4: p.4–14.
- Daily, G.C. 1997. *Nature's services: Societal Dependence on Natural Ecosystems*. Washington DC: Island Press.
- Dean, A.J., Barnett, A.G., Wilson, K.A., & Turrell, G. 2019. Beyond the 'extinction of experience' – Novel pathways between nature experience and support for nature conservation. *Global Environmental Change* 55: p.48–57. Available at: <http://www.sciencedirect.com/science/article/pii/S0959378018311336>.
- Drozdowski, I., Mrkvicka, A., & Pfundner, G. 2018. *Wiesen und Weiden im Wienerwald* 5th ed. Biosphärenpark Wienerwald Management GmbH.
- Egoh, B., Drakou, E., Dunbar, M., Maes, J., & Willemen, L. 2012. *Indicators for mapping ecosystem services: a review: JRC Scientific And Policy Reports*. Available at: http://bookshop.europa.eu/en/indicators-for-mapping-ecosystem-services-pbLBNA25456/;pgid=y8dlS7GUWMdSR0EAlMEUUsWb0000bkrukNw9;sid=WLTbRkQaKC9BsPRgnh_fC8Y_-ng9_Ol5Rsk=?CatalogCategoryID=7QwKABstDHwAAAEjK5EY4e5L.
- Ekins, P. 2011. Environmental sustainability: From environmental valuation to the sustainability gap. *Progress in Physical Geography* 35(5): p.629–651.
- European Commission. 2011. *Our life insurance, Our Natural Capital: An EU Bio-diversity Strategy to 2020*. Brussels: European Commission. Available at: <http://ec.europa.eu/environment/nature/biodiversity/comm2006/2020.htm>.
- Farber, S.C., Costanza, R., & Wilson, M.A. 2002. Economic and Ecological Concepts for Valuing Ecosystem Services. *Ecological Economics* 41(3): p.375–392.
- Fischer-Kowalski, M., & Erb, K.-H. 2004. Extending BRIM to BRIA. Social Monitoring and Integrated Sustainability Assessment. In C. Lee & T. Schaaf (eds) *Global Environmental and Social Monitoring*, 138–148. UNESCO
- Forman, R., & Gordon, M. 1986. *Landscape Ecology*. Wiley.
- Forman, R.T.T. 1995. *Land Mosaics: The Ecology of Landscapes and Regions*. Cambridge, UK: Cambridge University Press.
- von Geibler, J., Kristof, K., & Bienge, K. 2010. *Sustainability assessment of entire forest value chains: Integrating stakeholder perspectives and indicators in decision support tools*.

- Geofabrik GmbH, & OpenStreetMap Contributors. 2019. Download OpenStreetMap data for this region: Austria. Available at: <https://download.geofabrik.de/europe/austria.html> [Accessed March 2, 2019].
- Google Earth Pro. Wienerwald Biosphere Reserve Austria. 48° 08' 08.80"N, 16° 05'58.77"O Eye alt 54.32km. Image ©2019 DigitalGlobe ©2018 Google Image Landsat /Copernicus. Available at: <http://www.earth.google.com> [Accessed March 19, 2019].
- Grabherr, G., Koch, G., Kirchmeier, H., & Reiter, K. 1998. *Hemerobie österreichischer Waldökosysteme. Veröffentlichung des Österreichischen MAB-Programmes*. Innsbruck: Universitätsverlag Wagner.
- de Groot, R., Wilson, M., & Boumans, R. 2002. A typology for the classification, description and valuation of ecosystem functions, goods and services. *Ecological Economics* 41(May): p.1–20.
- de Groot, R.S. 1987. Environmental Functions as a Unifying Concept for Ecology and Economics. *The Environmentalist* 7(2): p.105–109.
- de Groot, R.S. 2006. Function-analysis and valuation as a tool to assess land use conflicts in planning for sustainable, multi-functional landscapes. *Landscape and Urban Planning* (75): p.175–186.
- de Groot, R.S., Alkemade, R., Braat, L., Hein, L., & Willemen, L. 2010. Challenges in integrating the concept of ecosystem services and values in landscape planning, management and decision making. *Ecological Complexity* 7(3): p.260–272.
- Haines-Young, & Potschin. 2010. The links between biodiversity, ecosystem service and human well-being. In D. G. Raffaelli & C. L. Frid (eds) *Ecosystem Ecology: A New Synthesis, BES Ecological Reviews Series*, Cambridge University Press
- Haines-Young, R., & Potschin, M. 2018. Common International Classification of Ecosystem Services (CICES) V5.1 and Guidance on the Application of the Revised Structure. *European Environment Agency*.
- Hainz-Renetzeder, C., Schneidergruber, A., Kuttner, M., & Wrabka, T. 2014. Assessing the potential supply of landscape services to support ecological restoration of degraded landscapes: A case study in the Austrian-Hungarian trans-boundary region of Lake Neusiedl. *Ecological Modelling* 295: p.196–206. Available at: <http://dx.doi.org/10.1016/j.ecolmodel.2014.07.001>.
- Heal, G. 2000. Valuing Ecosystem Services. *Ecosystems* 3(1): p.24–30.
- Hein, L., Van Koppen, K.K., de Groot, R.S., & Van Ierland, E.C. 2006. Spatial scales, stakeholders and the valuation of ecosystem services. *Ecological Economics* 57(2): p.209–228.
- Hermann, A. et al. 2014. Assessment framework for landscape services in European cultural landscapes: An Austrian Hungarian case study. *Ecological Indicators* 37(PART A): p.229–240. Available at: <http://dx.doi.org/10.1016/j.ecolind.2013.01.019>.
- Hermann, A., Schleifer, S., & Wrabka, T. 2011. The Concept of Ecosystem Services Regarding Landscape Research: A Review Imprint / Terms of Use. : p.1–37. Available at: [file:///C:/Users/joan/Documents/Mendeley Desktop/Hermann, Schleifer, Wrabka - 2011 - The Concept of Ecosystem Services Regarding Landscape Research A Review Imprint Terms of Use.pdf](file:///C:/Users/joan/Documents/Mendeley%20Desktop/Hermann,%20Schleifer,%20Wrabka%20-%202011%20-%20The%20Concept%20of%20Ecosystem%20Services%20Regarding%20Landscape%20Research%20A%20Review%20Imprint%20Terms%20of%20Use.pdf).
- Hou, Y., Burkhard, B., & Müller, F. 2013. Uncertainties in landscape analysis and ecosystem service assessment. *Journal of Environmental Management* 127: p.S117–S131. Available at: <http://dx.doi.org/10.1016/j.jenvman.2012.12.002>.
- Hughes, J., Richardson, M., & Lumber, R. 2018. Evaluating connection to nature and the relationship with conservation behaviour in children. *Journal for Nature Conservation* 45: p.11–19. Available

- at: <http://www.sciencedirect.com/science/article/pii/S1617138118300268>.
- Isbell, F., Tilman, D., Polasky, S., & Loreau, M. 2015. The biodiversity-dependent ecosystem service debt. *Ecology Letters* 18(2): p.119–134. Available at: <https://doi.org/10.1111/ele.12393>.
- Jacobs, S., Burkhard, B., Van Daele, T., Staes, J., & Schneiders, A. 2015. “The Matrix Reloaded”: A review of expert knowledge use for mapping ecosystem services. *Ecological Modelling*.
- Kaiser, G. et al. 2013. *Mapping tsunami impacts on land cover and related ecosystem service supply in Phang Nga, Thailand*.
- Kandziora, M., Burkhard, B., & Müller, F. 2013. *Mapping provisioning ecosystem services at the local scale using data of varying spatial and temporal resolution*.
- Kaule, G. 1986. *Arten- und Biotopschutz*. Ulmer.
- Konkoly-Gyuró, É. 2011. Definition of landscape services. In Wrba T. et al. (ed) *Biodiversity and Ecosystem Services as Scientific Foundation for the Sustainable Implementation of the Redesigned Biosphere Reserve ‘Neusiedler See’*. *Biological Sciences*., Available at: <http://dx.doi.org/10.1553/bioserv-neusiedler-sees>.
- Kuttner, M., Hainz-Renetzeder, C., Hermann, A., & Wrba, T. 2013. Borders without barriers - Structural functionality and green infrastructure in the Austrian-Hungarian transboundary region of Lake Neusiedl. *Ecological Indicators* 31: p.59–72. Available at: <http://dx.doi.org/10.1016/j.ecolind.2012.04.014>.
- Layke, C., Mapendembe, A., Brown, C., Walpole, M., & Winn, J. 2012. Indicators from the global and sub-global Millennium Ecosystem Assessments: An analysis and next steps. *Ecological Indicators* 17: p.77–87.
- Levrel, H., & Bouamrane, M. 2008. Instrumental Learning and Sustainability Indicators: Outputs from Co-Construction Experiments in West African Biosphere Reserves. *Ecology and Society* 13(1).
- Linehan, J.R., & Gross, M. 1998. Back to the future, back to basics: The social ecology of landscapes and the future of landscape planning. *Landscape and Urban Planning* (43): p.207–223.
- Loiskandl, G. 2005. Euromab - Plenary Presentation and Workshop Results. In *Presentation of the Wienerwald Biosphere Reserve and introduction to field trip*, 132–142.
- Lotze-Campen, H., Reusswig, F., & Stoll-Kleemann, S. 2008. Socio-Ecological Monitoring of Biodiversity Change. Building upon the World Network of Biosphere Reserves. *GAIA Ökologische Perspektiven für Wissenschaft und Gesellschaft* 17(S1): p.107–115.
- Lucht, W., & Pachauri, R. 2004. The Mental Component of the Earth System. *Earth System Analysis for Sustainability*: p.341–365. Available at: <http://ieeexplore.ieee.org/document/6300053>.
- Luck, G.W. et al. 2009. Quantifying the Contribution of Organisms to the Provision of Ecosystem Services. 59(3): p.223–235.
- Maes, J. et al. 2016. An indicator framework for assessing ecosystem services in support of the EU Biodiversity Strategy to 2020. *Ecosystem Services* 17: p.14–23.
- Martínez-Harms, M.J., & Balvanera, P. 2012. Methods for mapping ecosystem service supply: a review. *International Journal of Biodiversity Science, Ecosystem Services & Management* 8(1–2): p.17–25. Available at: <https://doi.org/10.1080/21513732.2012.663792>.
- MEA. 2005. *Millennium Ecosystem Assessment: Ecosystems and Human Well-Being: Synthesis*. Washington DC: Island Press. Available at: <https://uwe.rl.talis.com/link?url=https%3A%2F%2Fwww.millenniumassessment.org%2Fdocum>

ents%2Fdocument.354.aspx.pdf&sig=e4e5c0210a809ac41a5afd39d66c38ca0a9d59c5a6e9073de5183059559a51df.

- Müller, F. et al. 2008. Landscape approaches to assess environmental security: summary, conclusions, and recommendations. In I. Petrosillo et al. (eds) *Use of Landscape Sciences for the Assessment of Environmental Security*, 475–486. Dordrecht: Springer Netherlands
- Müller, F., & Burkhard, B. 2007. An ecosystem based framework to link landscape structures, functions and services. In Ü. Mander, H. Wiggering, & K. Helming (eds) *Multifunctional Land Use - Meeting Future Demands for Landscape Goods and Services*, 37–64. Springer
- Odling-Smee, L. 2005. Dollars and sense. *Nature* 437(7059): p.614–616. Available at: <https://doi.org/10.1038/437614a>.
- Opdam, P., Steingröver, E., & Rooij, S. Van. 2006. Ecological networks: A spatial concept for multi-actor planning of sustainable landscapes. *Landscape and Urban Planning* 75(3–4): p.322–332.
- OpenStreetMap Austria. 2019. Mitmachen – Mapper werden. Available at: <https://www.openstreetmap.at/mitmachen/> [Accessed March 6, 2019].
- van Oudenhoven, A.P., Petz, K., Alkemade, R., de Groot, R.S., & Hein, L. 2012. Indicators for assessing effects of management on ecosystem services. *Ecological Indicators* 21: p.110–122.
- Perrings, C., Duraiappah, A., Larigauderie, A., & Mooney, H. 2011. The biodiversity and ecosystem services science-policy interface. *Science* 331(6021): p.1139–1140.
- Pope, J., Annandale, D., & Morrison-Saunders, A. 2004. Conceptualising sustainability assessment. *Environmental Impact Assessment Review* (24): p.595–616.
- Potschin, M., & Haines-Young, R. 2011a. Ecosystem services: Exploring a geographical perspective. *Progress in Physical Geography* 35(5): p.575–594.
- Potschin, M., & Haines-Young, R. 2011b. Introduction to the Special Issue: Ecosystem Services. *Progress in Physical Geography* 35(5): p.571–574.
- Prévo, A.-C., Cheval, H., Raymond, R., & Cosquer, A. 2018. Routine experiences of nature in cities can increase personal commitment toward biodiversity conservation. *Biological Conservation* 226: p.1–8. Available at: <http://www.sciencedirect.com/science/article/pii/S0006320717319808>.
- Price, M.F., Gurung, A.B., Dourojeanni, P., & Maselli, D. 2006. Social Monitoring in Mountain Biosphere Reserves. *Mountain Research and Development* 26(2): p.174–180.
- Reimoser, F. et al. 2008. *ISWI - MAB Integrated Sustainable Wildlife Management in the Biosphere Reserve Wienerwald Integriertes nachhaltiges Wildtiermanagement im Biosphärenpark Wienerwald*. Wien.
- Reimoser, F. et al. 2012. *IESP – Grundlagen für eine integrierte ökologisch- räumliche Planung im Biosphärenpark Wienerwald. nachhaltiges Wildtiermanagement und Freizeitaktivitäten*.
- Ruoss, E. 2013. Biosphere Reserves as Model Sites for Sustainable Development . In M. Getzner & J. Michael (eds) *Protected Areas in Focus: Analysis and Evaluation. Proceedings in the Management of Protected Areas*, 99–114. Klagenfurt: Verlag Johannes Heyn
- Scolozzi, R., Morri, E., & Santolini, R. 2012. Delphi-based change assessment in ecosystem service values to support strategic spatial planning in Italian landscapes. *Ecological Indicators* 21: p.134–144.
- Seppelt, R. et al. 2012. Form follows function? Proposing a blueprint for ecosystem service assessments based on reviews and case studies. *Ecological Indicators* 21: p.145–154.

- Staudinger, M., Grass, V., Wrbka, T., Mair, B., & Rötzer, H. 2014. *Bericht Methodik der Projekte Offenlanderhebung Biosphärenpark Wienerwald 2011, Teil 1 und Offenlanderhebung Biosphärenpark Wienerwald 2012-2013, Teil 2*. Wien.
- Steffen, W. et al. 2011. The anthropocene: From global change to planetary stewardship. *Ambio* 40(7): p.739–761.
- Stoll, S. et al. 2015. Assessment of ecosystem integrity and service gradients across Europe using the LTER Europe network. *Ecological Modelling* 295: p.75–87. Available at: <http://dx.doi.org/10.1016/j.ecolmodel.2014.06.019>.
- Syrbe, R.U., & Walz, U. 2012. Spatial indicators for the assessment of ecosystem services: Providing, benefiting and connecting areas and landscape metrics. *Ecological Indicators* 21: p.80–88.
- TEEB ed. 2010. *TEEB - The Economics of Ecosystems and Biodiversity for Local and Regional Policy Makers*. Available at: <http://www.teebweb.org/ForLocalandRegionalPolicy/localandRegionalPolicyMakersChapterDrafts/tabid/29433/Default.aspx>.
- Termorshuizen, J.W., & Opdam, P. 2009. *Landscape services as a bridge between landscape ecology and sustainable development*.
- UNESCO. 1996. *Biosphere Reserves: The Seville Strategy and the Statutory Framework of the World Network*. Paris: UNESCO.
- UNESCO. 2002. *Biosphere reserves: Special places for people and nature*. Paris: UNESCO.
- UNESCO. 2005. *UNESCO and sustainable development*. Paris: UNESCO.
- UNESCO. 2016. Lima Action Plan for UNESCO 's Man and the Biosphere (MAB) Programme and its World Network of Biosphere Reserves (2016-2025). : p.1–11. Available at: http://www.unesco.org/fileadmin/MULTIMEDIA/HQ/SC/pdf/Lima_Action_Plan_en_final.pdf.
- United Nations. 2003. SEEA System of Environmental Economic Accounting. Available at: <https://seea.un.org/> [Accessed June 20, 2019].
- Vihervaara, P., Kumpula, T., Ruokolainen, A., Tanskanen, A., & Burkhard, B. 2012. Using detailed biotope data for Ecosystem service assessment in Natural protection areas. *International Journal of Biodiversity Science, Ecosystem Services & Management* 8(1–2): p.169–185.
- Vihervaara, P., Kumpula, T., Tanskanen, A., & Burkhard, B. 2010. Ecosystem services—A tool for sustainable management of human–environment systems. Case study Finnish Forest Lapland. *Ecological Complexity* 7(3): p.410–420.
- WCED. 1987. Our Common Future: Report of the World Commission on Environment and Development (Brundtland Report). In *World Commission on Environment and Development*, Oxford: Oxford University Press
- Weichhart, P. 2006. Raumbezogene Identität als Problemstellung der Regionalentwicklung. In *Beiträge zur theoretischen Grundlegung der Raumentwicklung*, 51–68. Hannover
- Westman, W.E. 1977. How Much Are Nature Services Worth? *Science* 197(4307): p.960–964. Available at: <http://science.sciencemag.org/content/197/4307/960.abstract>.
- Wienerwald Tourismus GmbH. 2018. Karte. Available at: <https://www.wienerwald.info/karte#&latitude=48.167665&longitude=16.110151999999997> [Accessed August 20, 2018].
- Willemen, L., Hein, L., van Mensvoort, M.E.F., & Verburg, P.H. 2010. Space for people, plants, and

livestock? Quantifying interactions among multiple landscape functions in a Dutch rural region. *Ecological Indicators* 10(1): p.62–73. Available at: <http://www.sciencedirect.com/science/article/pii/S1470160X09001472>.

Willemsen, L., Veldkamp, A., Verburg, P.H., Hein, L., & Leemans, R. 2012. A multi-scale modelling approach for analysing landscape service dynamics. *Journal of Environmental Management* 100: p.86–95. Available at: <http://www.sciencedirect.com/science/article/pii/S0301479712000308>.

Wilson, M.A., & Howarth, R.B. 2002. Discourse-based valuation of ecosystem services: establishing fair outcomes through group deliberation. *Ecological Economics* 41(3): p.431–443.

Wrbka, T. et al. 2012. *Man & Biosphere - Biodiversity and Ecosystem services as scientific foundation for the sustainable implementation of the Redesigned Biosphere Reserve "Neusiedler See" Final Report*.

9 Attachments

9.1 Definition of landscape services

Table A 1: BRIMSEN's definitions of the landscape services and correspondence with CICES V5.1 (Haines-Young et al. 2018). The examples are the authors' adaptation of de Groot (2006); de Groot, Wilson, and Boumans (2002) and Haines-Young and Potschin (2018).

Service Category	Code	Service	Definition	Simple descriptor	Examples Goods and Benefits
Regulation Services	RS1	Local climate and air regulation	Maintenance of a favourable local climate (temperature, humidity, precipitation, radiation) and air regulation by filtering wastes	Filtering wastes; Regulating the physical quality of air for people	Favourable local climate; Filtering effects from trees
	RS2	Disturbance prevention	Mitigation of environmental disturbances (flood prevention/mitigation; storm protection; wind protection; fire protection)	Protecting people from winds; Protecting people from fire; Stopping landslides and avalanches harming people; Regulating the flows of water in our environment	Enhanced safety from natural extreme events
	RS3	Water regulation	Regulation of runoff and river discharge	Regulating the flows of water in our environment	Drainage and natural irrigation
	RS4	Water supply	Filtering, retention and storage of fresh water for drinking, irrigation and industrial use	Controlling the chemical quality of freshwater	Pure water for drinking
	RS5	Soil retention	Role of vegetation root matrix and soil biota in holding soil (mitigation of erosion)	Controlling or preventing soil loss	Mitigated effects of erosion
	RS6	Soil formation	Weathering of rock, accumulation of organic matter	Ensuring soils form and develop	Maintenance of natural productive soils
	RS7	Nutrient regulation	Role of biota in storage and recycling of nutrients (e.g. N, P and S)	Ensuring the organic matter in our soils is maintained; Controlling the chemical quality of freshwater	Maintenance of healthy and productive ecosystems
	RS8	Pollination	Role of biota in movement of floral gametes	Pollinating our fruit trees and other plants	Edible fruits
Habitat Services	HS1	Refugium	Suitable living space for wild plants and animals	Providing habitats for wild plants and animals that can be useful to us	Forest edge for rodents; church towers for falcons
	HS2	Nursery	Suitable reproduction habitat	Providing habitats for wild plants and animals that can be useful to us	Extensive meadow for bird breeding

Provisioning Services	PS1	Food	Conversion of solar energy into wild and cultivated edible plants and animals	Any crops and fruits grown by humans for food; food crops; Plants that are cultivated in fresh or salt water that we eat; Livestock raised in housing and/or grazed outdoors; Animals that are cultivated in fresh or salt water that we eat; Food from wild plants, Food from wild animals	Edible wild plants, crops, fungi, livestock and fish
	PS2	Raw materials	Conversion of solar energy into biomass	Material from plants, fungi, algae or bacterial that we can use; Material from animals that we can use; Plant materials used as a source of energy; Materials from wild plants; Materials from wild animals; Seed collection	Material for human construction (building, roofing material and manufacturing) like fuel and energy wood; sand and gravel;
	PS3	Genetic resources	Genetic material and evolution in wild plants and animals	Seed collection; Plants, fungi or algae that we can use for breeding; Genetic material from wild plants, fungi or algae that we can use; Animals used for replenishing stock; Wild animals that we can use for breeding; The genetic information that is stored in wild animals that we can use	Improve crop quality
	PS4	Medicinal resources	Variety in chemical substances in natural biota	Materials from wild animals and from wild plants for non-nutritional uses	Drugs and pharmaceuticals
Carrier Services	CS1	Cultivation	Providing suitable substrate for cultivation	Any crops and fruits grown by humans for food; food crops;	Cultivated food and fodder
	CS2	Waste disposal	Providing suitable substrate for waste (including wastewater)	Decomposing wastes; filtering wastes	Space for waste disposal
	CS3	Habitation	Providing suitable space for human living	N/A	Living space
	CS4	Energy conversion	Providing suitable substrate or medium for energy conversion	N/A	Energy facilities
	CS5	Transportation	Providing suitable substrate or medium for transportation	N/A	Main and side roads as well as railroad tracks
	CS6	Tourism facilities	Providing space and facilities for human activities related to tourism	N/A	Accommodation for touristic use
Information Services	IS1	Aesthetic information	Attractive landscape features and views	The beauty of nature	Appreciation of blooming trees or meadows
	IS2	Recreation	Variety in landscapes with (potential) recreational uses	Using the environment for sport and recreation; using nature to help stay fit; Watching plants and animals where they live; using nature to destress;	Use of meadows for leisure activities "Lagerwiesen"

IS3	Cultural and artistic information	Variety in natural and cultural features with cultural and artistic value	the things in nature used to make films or to write books; The things in nature that help people identify with the history or culture of where they live or come from, Using nature to as a national or local emblem	Use of characteristic tree rows or single trees of cultural landscape in artistic display
IS4	Spiritual and historic information	Variety in natural and cultural features with spiritual and historical value	The things in nature that help people identify with the history or culture of where they live or come from; Using nature to as a national or local emblem, The things in nature that have spiritual importance for people	Provision of religious and historic elements like churches and crucifixes
IS5	Science and education	Variety in nature with scientific and educational value	Researching nature; studying nature	Possibility to experience, observe and learn about the environment e.g. species diversity, educational trails

9.2 Additional Spatial Indicators sources

Table A 2: Data sources of the Additional Spatial Indicators (ASI) classes, and the ASI types included in each class. The codes in brackets in the ASI types refer to the © OpenStreetMap 2018.

No	Code	ASI Classes	ASI Types
1	HYP1	Hydro power plants*	Hydro power plants
2	PUT2	Public transport	OSM: railways, bus stops
3	ROD3	Roads primary	OSM: Motorway, trunk, primary + all links
4	ROD4	Roads secondary	OSM: secondary, tertiary + all links
5	ROD5	Roads tertiary	OSM: unclassified
6	ROD6	Roads residential	OSM: residential, service roads
7	RTR7	Recreational trails	OSM: Bridleways; WWTourism: Nordic walking/jogging trails; Hiking - OSM: track all grades, path, footway, steps; WWTourism: Hiking trails, Cycling - OSM: cycle ways - paved; WWTourism: cycling/mtb trails
8	ETR8	Educational trails	WWTourism: `Themenwege`
9	REL9	Religious buildings	OSM: churches and chapels (3100+3102+3107)
10	REL10	Religious elements	OSM: wayside cross/shrine (2734+2735)
11	ART11	Monuments and artwork	OSM: memorial (2724), monument (2723), artwork (2725), fountain (2904)
12	ARC12	Archaeological sites	OSM: archaeological sites (2733)
13	MUS13	Museums	OSM: museum (2722)
14	CAS14	Castles	OSM: castles (2731), ruins (2732)
15	VPS15	View points	OSM: observation tower (2953), viewpoint (2742)
16	INF16	Recreational infrastructure	OSM: dog park (2206), playgrounds (2205), picnic site (2741), toilet (2901), drinking water facilities (2903), waste baskets (2906), bench (2902), tourist info (2701)
17	GST17	Gastronomy	OSM: biergarten (2307), restaurant (2301), cafe (2303), pub (2304), fast-food (2302), Bakery (2502), bar (2305)
18	ACC18	Accommodation	OSM: hotel (2401), guesthouse (2404), hostel (2405), campsite (2422), alpine hut (2423)

OSM: © OpenStreetMap contributors 2018. download 2.3.2019:

<https://download.geofabrik.de/europe/austria.html>

WWTourism: Wienerwald Tourismus GmbH 2018. download (gpx) 20.8.2018:

<https://www.wienerwald.info/karte>

*data dismissed (with landscape services Energy conversion) Source: Institute of Hydrobiology and Aquatic Ecosystem Management (IHG) of the University of Natural Resources and Life Sciences

9.3 Reclassification of open land elements

Table A 3: Reclassification of the open land types (OL types) into open land classes (OL classes) of the Wienerwald Biosphere Reserve. OL types refer to the open land types available in Staudinger et al. (2014)

No	Code	OL Class	OL Type	Notes
1	WT1	Straight, semi natural streams	2	
2	WT2	Meandering, semi natural streams	3, 4, 19	includes submerged vegetation in streams (19)
3	WT3	Channel/regulated streams	7,8, 133	includes straight lowland streams (133)
4	WT4	Periodic/small brooks	9,10	includes ditches (9)
5	WT5	Semi natural, nutrient-poor lakes and pools	11,12,14, 18	includes (meso) eutrophic pools (12), episodic pools (14), submerged vegetation in standing water (18)
6	WT6	Nutrient-rich pools	13,15	includes nutrient-rich not natural pools connected to groundwater (15)
7	WT7	Artificial standing water connected to groundwater	17	
8	WT8	Artificial pools disconnected from groundwater	16	includes species-poor eutrophic pools
9	WT9	Springs, marshes, swamps	21,23	
10	WT10	Nutrient-poor fens	24,25	includes acidophilic and basophilic fens
11	WT11	Reed beds	26,27,28,29,31	
12	TH12	Nutrient-rich moist tall herbs	30,33,34,35, 36,44,45,46, 47,48,58	includes ruderal reed beds (30), corridors dominated by <i>Petasites hybridus</i> (33), corridors dominated by <i>Filipendula hydrophilous</i> (34), wet abandoned pastures dominated by umbellifers (35), corridors dominated by <i>Senecio sarracenicus</i> (36), moist fallow land dominated by <i>Petasites hybridus</i> (44), moist fallow land dominated by <i>Filipendula hydrophilous</i> (45), moist fallow land dominated by umbellifers (46), abandoned reed beds (47), abandoned nutrient- rich wetlands (48), fallow land of damp (semi-moist) grasslands (58)
13	MD13	Nutrient-poor, extensive moist meadow	37,38,40,42,55	includes straw meadow dominated by <i>Molinia caerulea</i> (37), nutrient-poor fallow land of damp (semi-moist) and wet grasslands (38), nutrient-poor <i>Molinia</i> meadows (40), unfertilized marshy meadows (42), semi-humid oat grass meadows (<i>Filipendulo vulgaris</i> - <i>Arrhenatheretum</i>). It is part of EU habitat type: 6410 <i>Molinia</i> meadows on calcareous, peaty or clayey-silt-laden soils, mire meadows, 6510 lowland hay meadows.

14	MD14	Nutrient-rich moist and wet meadow	41,43, 207	includes fertilized moist and wet meadows. Typical species: <i>Cirsium rivulare</i> , <i>Holcus lanatus</i> , <i>Ranunculus acris</i> , <i>Festuca pratensis</i> , <i>Lychnis flos-cucculi</i> , <i>Carex panicea</i> (41); moist and wet pastures (43), regularly flooded prairies (207)
15	SB15	Nutrient-poor moist shrub succession	39	Secondary succession of fallow land of moist meadows
16	SB16	Nutrient-rich moist shrub succession	49, 108	Fallow Land of nutrient-rich moist meadows (49), shrub succession (108)
17	SB17	Shrubs on semi-moist grassland	59	shrubs on damp (moderate covering of moisture) grassland
18	TR18	Chasm woodland	132,109	chasm woods (109) and abandoned fruit trees with shrub succession (132)
19	TR19	Abandoned fruit trees	140	
20	TH20	Nutrient-rich tall herbs	50,52	includes Nettle meadows, neophytes like <i>Impatiens glandulifera</i> and <i>Reynoutria</i> spp.
21	TH21	Nutrient-rich herbs (blooming aspect)	51	<i>Solidago</i> sp. (Golderod)
22	MD22	Semidry managed meadows	54,74,75,76	includes dry oat grass land (54), dry brome grass (74), semi-dry brome grass (75), secondary grazed semi-dry pastures (76). It is part of EU habitat type: 6510 Lowland hay meadows, 6210 Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia, important orchid sites)
23	MD23	Intermediate managed meadows	56,57,143,204	includes oat grassland (56), damp meadows dominated by <i>Alopecurus</i> (57), grazed Pastinaco-Arrhenatheretum (143), mountainous Astrantio-Trisetetum (204). It is part of EU habitat type: 6510 Lowland hay meadows, rich pastures, 6520 Mountain hay meadows
24	MD24	Intensively managed meadows and pastures	60,63,67	includes intensively managed pastures (60)(67), field forage (63)
25	DR25	Semi-dry and dry fallow land of fields and vineyards	61,62,77	includes fallow land of semi-dry and dry grassland (77)
26	MD26	Nutrient-poor managed meadows and pastures	64,65,66	includes acidophilic (65) and basophilic (66) pastures, Red Fescue meadows (64)
27	DR27	Dry shrubland on limestone	69	
28	DR28	Thermophilous dry shrubland	70	EU habitat type: 40A0 Subcontinental peri-pannonic scrub
29	DR29	Semidry and dry grassland	72,73,78, 134	includes rupicolous dry grasslands (72), steppic grasslands (73), fallow land of semi-dry and dry grasslands (78), 6190 Rupicolous pannonic grasslands (Stipo-Festucetalia pallentis), 6240 Sub-Pannonic steppic grasslands, 6210 Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia, * important orchid sites), thermophilous scree vegetation (134)

30	DR30	Dry forest edges	79	
31	TR31	Forest edges	0_Wald, 135,136,199, 210-228	
32	TR32	<i>Pinus nigra</i> forests on dry grassland	200, 202	includes natural <i>Pinus nigra</i> stands in open land (202)
33	MG33	Species-rich slopes and margins	82,83	includes field margins with pannonic shrub fringes (83)
34	RC34	Rock fragments and dry-stone walls	113	
35	MG35	Species-poor field margins	84,85	includes ruderal species (84,85)
36	MG36	Ruderal Vegetation	80, 86	
37	CU37	Cultivated field	87	
38	CU38	Vineyard	88	
39	CU39	Intensively managed orchard	105	
40	SB40	Nitrophilous and neophytic edges and woodlets	89,90,93	
41	SB41	Hedges and shrubs	91	
42	SB42	Shrubs on moist soil	92	
43	TR43	Long-rotation woodland	94,96,103	includes <i>Robinia pseudoacacia</i> (94), poplar, birch, spruce woodlands, non-natural hedges and windbreaks (96), copses of allochthonous species (103)
44	TR44	Hedgerows dominated by trees	95,102	includes hedgerows (95), native and common deciduous trees (maple, cherry tree, linden, etc.) (102)
45	TR45	Tree rows and single trees	97,101	
46	TR46	Larch dominated meadows/pastures	205	
47	TR47	Riparian woodland	98,99,100	includes 91F0 Riparian mixed forests of <i>Quercus robur</i> , <i>Ulmus laevis</i> and <i>Ulmus minor</i> , <i>Fraxinus excelsior</i> or <i>Fraxinus angustifolia</i> , along the great rivers (<i>Ulmion minoris</i>), non-natural riparian woodland
48	TR48	Extensive orchards	104	
49	TR49	Short-rotation plantations	106,107	includes "energy forest", tree nurseries
50	RC50	Cliffs	112	
51	RC51	Active gravel pits and quarries	115,116	
52	RC52	Abandoned stone quarries	117	
53	RC53	Abandoned gravel pits	118	
54	AR54	Leisure and sport facilities	119	includes turf
55	AR55	Lawns	123	
56	GD56	Gardens or parks	120, 121	
57	GD57	Greenery and planting	122	

58	GD58	Cemeteries	124	
59	AR59	Unvegetated open areas with compacted soil including parking areas, storage sites	125	
60	AR60	Paved open areas including parking areas, storage sites	126	
61	AR61	Isolated buildings	127	
62	AR62	Waste sites	129	includes wastewater treatment sites

9.4 Classification of Additional Spatial Indicators

Table A 4: Classification of Additional Spatial Indicators (ASI) for the Wienerwald Biosphere Reserve.

No	Code	ASI Class	Description
1	HYP1	Hydro Power Plants*	
2	PUT2	Public transport	Railway lines and bus stops providing public transportation
3	ROD3	Roads primary	Major roads, including highways and roads of national reach/extent
4	ROD4	Roads secondary	Major roads of regional or local extent
5	ROD5	Roads tertiary	Small local roads
6	ROD6	Roads residential	Roads in residential areas providing access to buildings
7	RTR7	Recreational trails	Hiking: footways, paths and tracks for pedestrian use (including unpaved/gravel roads of forestry and agriculture). Cycling: paths paved/unpaved for cycling and mountain biking. Nordic Walking/Jogging: paths used for jogging and Nordic walking. Bridleways: paths used for horse riding
8	ETR8	Educational trails	Trails providing information or interactive education
9	REL9	Religious buildings	Sacral buildings like churches and chapels
10	REL10	Religious elements	Small sacral elements including wayside crucifixes and shrines
11	ART11	Monuments and artwork	Built elements including memorials, monuments, fountains and artwork.
12	ARC12	Archaeological sites	Archaeological excavations
13	MUS13	Museums	
14	CAS14	Castles and ruins	
15	VPS15	View points	Overlooks providing view of the landscape, including lookout-towers
16	INF16	Recreational Infrastructure	Built and maintained elements and facilities providing recreational infrastructure including playground, picnic area, dog areas, benches, drinking water, waste baskets, toilets
17	GST17	Gastronomy	Facilities in the gastronomical sector e.g. restaurant, café, pub, bakery, beer garden
18	ACC18	Accommodation	Touristic facilities providing accommodation e.g. hotel, guesthouse, hostel, campsite, alpine hut

*Data dismissed (with LS Energy conversion)

10 Teaching Landscape Services in GIS - three case studies

The concept of landscape and ecosystem services is still rather new in academic teaching. Most of landscape service-related research deals with service mapping. For this reason, it is very relevant to include it in GIS courses.

In this chapter, three case studies, which made use of some BRIMSEN sources, are presented. Although the following three case studies are associated to the project, a different data base was used (including forest and the SE region of the Biosphere Reserve); therefore, the results obtained through the course are not comparable to those obtained in BRIMSEN.

The University of Natural Resources and Life Sciences offers regularly a lecture called „GIS-gestütztes Projekt“ with the aim to teach applied GIS-competences to Bachelor students of the curriculum „Landschaftsplanung und Landschaftsarchitektur“ (landscape planning and landscape architecture). The lecturer has the possibility to implement own ideas or project-based research questions in this course. Also, the students themselves are invited to formulate concrete research questions within this frame. In all, the work load of this lecture is five ECTS points, which equals 125 hours of work for the student. In the winter semester 2018/19 one course was dedicated to questions related to the BRIMSEN project. In order to facilitate questions related to landscape services provided by the Wienerwald Biosphere Reserve, focus was on three of the five service groups (regulation, habitat, provision services) where land use was the only used indicator in the capacity matrix and no additional spatial indicators had to be used. Three groups of students were established, each looking at a different topic related to landscape services in the Wienerwald Biosphere Reserve.

In conclusion, the concept of landscape services was well understandable for the students although implementing in GIS presented a challenge. The details of what exactly is calculated and what do the results mean was hard to grasp for them.

Team A: Aggregation of landscape services to service groups and on district level *by Lena Graf and Eric Kraus; Supervisor: Christa Hainz-Renetzeder*

Landscape Service Maps on different scale with different resolution are widely available. But quantification into numbers still provides a challenge. Burkhard (2017) explicitly addresses major issues connected with the selection of relevant geo-biophysical spatial units and linking these with landscape services. We wanted to investigate aggregation procedures of landscape services in the Wienerwald Biosphere Reserve and to answer the following questions: (1) What is the average landscape service capacity in the individual political districts of the Wienerwald Biosphere Reserve? and (2) Is there a difference in including area-weighting into the averaging calculation?

For calculation of the average landscape service, we used two slightly different methods. First, we calculated the mean of each landscape service for each district and then averaged for each of the three service groups (regulation, habitation, provision). Second, we calculated the area of each landscape element of open land classes and its percentage of the whole area for each class in the district and then averaged over the landscape service in order to receive an area-weighted mean. This area-weighted mean was again averaged for each of the three service groups (for more details, see method section in Hermann et al., 2014).

Averaging the individual landscape services into the three main service groups for each

political district showed large differences among them (Table A.1). Looking at the values without area-weighting, provision services were always lowest. Regulation and Habitat services were balanced throughout the districts. Including the size of the landscape elements of open land classes into the averaging procedure changed the picture immensely.

Table A.1: Mean capacity values for each landscape service groups per district; on the right including area-weights.

District	mean landscape service			area-weighted mean landscape service		
	regulation	habitat	provision	regulation	habitat	provision
Wien	3,29	3,43	2,12	2,36	1,94	1,30
Baden	2,78	2,74	1,62	3,22	1,97	1,28
Lilienfeld	2,78	2,58	1,56	3,54	3,62	2,38
Moedling	2,72	2,66	1,59	3,77	3,97	2,69
St. Poelten	2,81	2,72	1,71	3,48	3,68	2,59
Tulln	2,64	2,61	1,67	3,90	4,17	2,79

Averaging landscape service values into service groups and for spatial units is largely connected with information loss as already stated by Burkhard (2017). We argue that the size of the landscape elements (grouped by open land class) has a large impact on how much landscape service capacity is referenced to in each unit.

Team B: Landscape Services and Hemeroby by *Daniel Hartl and Tobias Riffel*; Supervisor: *Christa Hainz-Renetzeder*

The concept of hemeroby was originally developed for measuring human impacts on flora and vegetation. Later this concept was applied on whole ecosystems (Blume & Sukopp 1976: p. 83; Sukopp 1972: p. 113ff). The hemerobiotic state of an area is expressed as the magnitude of the deviation from the potential natural vegetation caused by human activities. The aim of this report is to look at the following aspect: do the open land classes with the same degree of hemeroby have similar capacity values of landscape services?

We assigned for each open land class (and forest) a degree of hemeroby and implemented this classification into GIS for the Wienerwald Biosphere Reserve. We looked at the distribution of the matrix values for the landscape services in each hemerobic class. Frequency analysis was performed for each landscape service in ArcMap 10.4.1 (ESRI, Redlands) and analysed in SPSS and Excel. There were nearly no ahemerobic spatial units (1.5%), but large areas are covered by oligohemerobic land-use elements (66.2 %; mainly forest). Also, polyhemerobic (4.9%) and metahemerobic (15.8%) polygons are present. Regulation services showed the trend that metahemerobic elements only showed minor values for their capacity and the highest capacity value 5 was nearly not present (Figure B.1.a as example for "Disturbance prevention"). Capacity values 3 and 4 were existent in all degrees of hemeroby, in some landscape services a peak of capacity value 3 and 4 can be seen in the oligohemerob category. High capacity values for Habitat services clearly were given in low hemerobic categories, whereas low capacity values were clustered in the metahemerobic category (Figure B.1.b as example for "Refugium"). Provision services in general showed only low capacity values, as such a trend for a distribution among the degrees of hemeroby could not be found. For rough assessments and if no detailed land cover classification is available, hemeroby values can help in estimating landscape service capacity especially for regulating and habitat services. For a local scale service assessment, a more detailed classification however is needed.

Figure B.1: a. Bar chart of landscape service “Disturbance prevention”; b. Bar chart of landscape service “Refugium”. The height of the bar is representing the count of each value in the capacity matrix, the colour of the bar is visualizing a certain capacity value.

Team C: Local Climate Regulation as an example for an important landscape service in and near urban areas by Michele Misztowt and Theresa Rockenschaub; Supervisor: Christa Hainz-Renetzedner

The term "urban heat island" describes built up urban areas, mostly insular, that are hotter than nearby rural areas (Hupfer & Kuttler, 2006). The annual mean air temperature of a Central European major city is 1°C warmer than its surroundings. Especially during the night, the difference can be as high as 10°C. Water bodies and forest counterbalance the negative effects of human-made areas like streets, vineyards, fields. A high share of sealed surfaces leads to reduced evaporation in nearby villages whereas insolation is saved in buildings and streets. Green infrastructure and green spaces can reduce heat storage, raise evaporation and thus lead to cooling (Mursch-Radlgruber, 2017). This report focuses on open land classes of the Biosphere Reserve and their capacity to regulate local climate. We present the distribution of capacity values for local climate regulation and compare this with climate data on the area.

In GIS, the landscape service capacity matrix was joined to open land classes and “local climate regulation” was selected for further calculations such as summary statistics.

In the Wienerwald Biosphere Reserve large areas showed a high capacity for local climate regulation which is mostly due to their big forest areas. Nevertheless, especially at the border of the biosphere reserve, many elements occur with low local climate regulation. Looking at the highest temperature in summertime (Fig. C.1.a) and wintertime (Fig. C.1.b.), a similar picture can be seen. The highest temperatures exist along the border of the biosphere reserve where the most elements with a low capacity can be found. Still, as temperature is strongly dependent on sea level (lapse rate), we also need to take account of this information. And as seen in Figure C.1, also a strong correlation with altitude is visible.

Figure C.1: a. on the left; highest mean temperature in August; b. on the right; highest mean temperature in January. Contour lines in grey, Numbers along contour lines means altitude.

References

- Blume, H.-P., & Sukopp, H. (1976): Ökologische Bedeutung anthropogener Bodenveränderungen, [Ecological significance of anthropogenic soil changes]. Schriftenreihe für Vegetationskunde, 10, 75–89.
- Burkhard, B. (2017): Ecosystem services matrix. In: Burkhard, B. & Maes, J. (Eds) Mapping Ecosystem Services. Pensoft Publishers, Sofia, 227-232.
- Fick, S.E., Hijmans, R.J. (2017): Worldclim 2: New 1-km spatial resolution climate surfaces for global land areas. International Journal of Climatology.
- Hermann, A., Kuttner, M., Hainz-Renetzeder, C., Konkoly-Gyuró, E., Tirászi, A., Brandenburg, C., Allex, B., Ziener, K., Wrbka, T. (2014): Assessment framework for landscape services in European cultural landscapes: An Austrian Hungarian case study; Ecological Indicators 37, 229-240.
- Hupfer P., Kuttler W. (2006): Witterung und Klima. Eine Einführung in die Meteorologie und Klimatologie. [Weather and Climate. An Introduction in Meteorology and Climatology.] 12. Aufl. Wiesbaden.
- Kowarik, I. (1988): Zum menschlichen Einfluss auf Flora und Vegetation: Theoretische Konzepte und ein Quantifizierungsansatz am Beispiel von Berlin (West), [To the human impact on flora and vegetation: Theoretical concepts and a quantification approach using the example of Berlin (West)]. Landschaftsentwicklung und Umweltforschung, 56, 1–280.
- Mursch-Stadlgruber, E. (2017): Stadt- und Geländeklimatologie [Urban and Terrain Climatology]. Vorlesung WiSe 2017 [Lecture Winterterm 2017]. Wien.
- Sukopp, H. (1976): Dynamik und Konstanz in der Flora der Bundesrepublik Deutschland, [Dynamics and stability in Flora of the Federal Republic of Germany]. Schriftenreihe für Vegetationskunde, (10), 9–26.